

Cullin 4A/4B Rabbit mAb

Catalog # AP75306

Product Information

Application	WB, IHC-P
Primary Accession	<u>Q13620</u>
Reactivity	Human, Rat, Hamster
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	103982

Additional Information

Gene ID	8450
Other Names	CUL4B
Dilution	WB~~1/500-1/1000 IHC-P~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CUL4B {ECO:0000303 PubMed:14578910, ECO:0000312 HGNC:HGNC:2555}
Function	 Core component of multiple cullin-RING-based E3 ubiquitin- protein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:<u>14578910</u>, PubMed:<u>16322693</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>, PubMed:<u>22118460</u>, PubMed:<u>29779948</u>, PubMed:<u>30166453</u>, PubMed:<u>33854232</u>, PubMed:<u>33854239</u>). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition subunit (PubMed:<u>14578910</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>, PubMed:<u>22118460</u>, PubMed:<u>29779948</u>). CUL4B may act within the complex as a scaffold protein, contributing to catalysis through positioning of the substrate and the ubiquitin- conjugating enzyme (PubMed:<u>14578910</u>, PubMed:<u>16678110</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>, PubMed:<u>14578910</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>, PubMed:<u>16678110</u>, PubMed:<u>18593899</u>). Targeted to UV damaged chromatin by DDB2 and may be important for DNA repair and DNA replication (PubMed:<u>16678110</u>). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component)

are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed: 29779948). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiquitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:33854232, PubMed:33854239). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed: 30166453). Required for ubiguitination of cyclin E (CCNE1 or CCNE2), and consequently, normal G1 cell cycle progression (PubMed: 16322693, PubMed: 19801544). Regulates the mammalian target-of- rapamycin (mTOR) pathway involved in control of cell growth, size and metabolism (PubMed:<u>18235224</u>). Specific CUL4B regulation of the mTORC1- mediated pathway is dependent upon 26S proteasome function and requires interaction between CUL4B and MLST8 (PubMed:<u>18235224</u>). With CUL4A, contributes to ribosome biogenesis (PubMed:26711351).

Cellular LocationCytoplasm {ECO:0000250 | UniProtKB:A2A432}. Nucleus. Note=More
concentrated in nuclei than in cytoplasm in germinal vesicle (GV) stage
oocytes, zygotes and the 2-cell stage, but distributed in the cytoplasm at the
MII-stage oocytes. {ECO:0000250 | UniProtKB:A2A432}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.