

DAPK2 Rabbit mAb

Catalog # AP75337

Product Information

Application	WB, IHC-P, FC
Primary Accession	Q9UIK4
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	42898

Additional Information

Gene ID	23604
Other Names	DAPK2
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A FC~~1:200-1:1000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	DAPK2
Function	Calcium/calmodulin-dependent serine/threonine kinase involved in multiple cellular signaling pathways that trigger cell survival, apoptosis, and autophagy. Regulates both type I apoptotic and type II autophagic cell death signals, depending on the cellular setting. The former is caspase-dependent, while the latter is caspase-independent and is characterized by the accumulation of autophagic vesicles. Acts as a mediator of anoikis and a suppressor of beta-catenin-dependent anchorage-independent growth of malignant epithelial cells. May play a role in granulocytic maturation (PubMed: 17347302). Regulates granulocytic motility by controlling cell spreading and polarization (PubMed: 24163421).
Cellular Location	Cytoplasm. Cytoplasmic vesicle, autophagosome lumen
Tissue Location	Expressed in neutrophils and eosinophils (PubMed: 24163421). Isoform 2 is expressed in embryonic stem cells (at protein level). Isoform 1 is ubiquitously

expressed in all tissue types examined with high levels in heart, lung and skeletal muscle



Background

Calcium/calmodulin-dependent serine/threonine kinase involved in multiple cellular signaling pathways that trigger cell survival, apoptosis, and autophagy. Regulates both type I apoptotic and type II autophagic cell death signals, depending on the cellular setting. The former is caspase-dependent, while the latter is caspase-independent and is characterized by the accumulation of autophagic vesicles. Acts as a mediator of anoikis and a suppressor of beta-catenin-dependent anchorage-independent growth of malignant epithelial cells. May play a role in granulocytic maturation (PubMed:17347302). Regulates granulocytic motility by controlling cell spreading and polarization (PubMed:24163421).

Images



Western blot analysis of DAPK2 in rat spleen lysates using DAPK2 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.