

[KD]DKC1 Rabbit mAb

Catalog # AP75361

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	O60832
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	57674

Additional Information

Gene ID	1736
Other Names	DKC1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IHC-F~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	DKC1 (HGNC:2890)
Synonyms	NOLA4
Function	[Isoform 1]: Catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA (PubMed: 25219674 , PubMed: 32554502). This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1 (PubMed: 25219674). Each rRNA can contain up to 100 pseudouridine ('psi') residues, which may serve to stabilize the conformation of rRNAs. Required for ribosome biogenesis and telomere maintenance (PubMed: 19179534 , PubMed: 25219674). Also required for correct processing or intranuclear trafficking of TERC, the RNA component of the telomerase reverse transcriptase (TERT) holoenzyme (PubMed: 19179534).
Cellular Location	[Isoform 1]: Nucleus, nucleolus. Nucleus, Cajal body {ECO:0000250 UniProtKB:P40615}


Tissue Location

Ubiquitously expressed.


Background

Required for ribosome biogenesis and telomere maintenance. Probable catalytic subunit of H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA. This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1.



Images



Western blot analysis of DKC1 in HeLa(WT) and DKC1-Knockdown HeLa(KD) using DKC1 antibody.




Immunocytochemistry analysis of DKC1 (green) in K562 using DKC1 antibody, and DAPI (blue).



Western blot analysis of DKC1 in K562, C6, 3T3, HeLa lysates using DKC1 antibody.

Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using DKC1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.