

# DNA Ligase IV Rabbit mAb

Catalog # AP75364

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">P49917</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	103971

## Additional Information

---

<b>Gene ID</b>	3981
<b>Other Names</b>	LIG4
<b>Dilution</b>	WB~~1:500-1:1000 IHC-P~~N/A
<b>Format</b>	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	LIG4 {ECO:0000303   PubMed:16357942, ECO:0000312   HGNC:HGNC:6601}
<b>Function</b>	DNA ligase involved in DNA non-homologous end joining (NHEJ); required for double-strand break (DSB) repair and V(D)J recombination (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">23523427</a> , PubMed: <a href="#">29980672</a> , PubMed: <a href="#">33586762</a> , PubMed: <a href="#">8798671</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). Catalyzes the NHEJ ligation step of the broken DNA during DSB repair by resealing the DNA breaks after the gap filling is completed (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). Joins single-strand breaks in a double-stranded polydeoxynucleotide in an ATP-dependent reaction (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). LIG4 is mechanistically flexible: it can ligate nicks as well as compatible DNA overhangs alone, while in the presence of XRCC4, it can ligate ends with 2-nucleotides (nt) microhomology and 1-nt gaps (PubMed: <a href="#">17290226</a> ). Forms a subcomplex with XRCC4; the LIG4-XRCC4 subcomplex is responsible for the NHEJ ligation step and XRCC4 enhances the joining activity of LIG4 (PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). Binding of the

LIG4-XRCC4 complex to DNA ends is dependent on the assembly of the DNA-dependent protein kinase complex DNA-PK to these DNA ends (PubMed:[10854421](#)). LIG4 regulates nuclear localization of XRCC4 (PubMed:[24984242](#)).


**Cellular Location** Nucleus

**Tissue Location** Testis, thymus, prostate and heart.



## Background

Efficiently joins single-strand breaks in a double-stranded polydeoxynucleotide in an ATP-dependent reaction. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination.

## Images



Western blot analysis of DNA Ligase IV in K562, U251MG, HepG2, MCF-7 lysates using DNA Ligase IV antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.