

DOCK2 Rabbit mAb

Catalog # AP75368

Product Information

Application WB, IHC-P
Primary Accession Q92608
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 211948

Additional Information

Gene ID 1794

Other Names DOCK2

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format Liquid

Protein Information

Name DOCK2

Synonyms KIAA0209

Function Involved in cytoskeletal rearrangements required for lymphocyte migration

in response of chemokines. Activates RAC1 and RAC2, but not CDC42, by functioning as a guanine nucleotide exchange factor (GEF), which exchanges bound GDP for free GTP. May also participate in IL2 transcriptional activation

via the activation of RAC2.

Cellular Location Endomembrane system; Peripheral membrane protein. Cytoplasm,


cytoskeleton. Note=Colocalizes with F-actin

Tissue Location Specifically expressed in hematopoietic cells. Highly expressed in peripheral

blood leukocytes, and expressed at intermediate level in thymus and spleen.

Expressed at very low level in the small intestine and colon.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.