

# Phospho-EGFR (Tyr1068) Rabbit mAb

Catalog # AP75378

## Product Information

|                   |                           |
|-------------------|---------------------------|
| Application       | WB, IHC-P, IHC-F, IF, ICC |
| Primary Accession | <a href="#">P00533</a>    |
| Reactivity        | Human                     |
| Host              | Rabbit                    |
| Clonality         | Monoclonal Antibody       |
| Calculated MW     | 134277                    |

## Additional Information

|             |   |
|-------------|---|
| Gene ID     | 1956  |
| Other Names | EGFR  |
| Dilution    | WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IF~~1:50~200 ICC~~N/A                          |
| Format      | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |

## Protein Information

|          |  |
|----------|--|
| Name     | EGFR ( <a href="#">HGNC:3236</a> )   |
| Synonyms | ERBB, ERBB1, HER1  |
| Function | <p>Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:<a href="#">10805725</a>, PubMed:<a href="#">27153536</a>, PubMed:<a href="#">2790960</a>, PubMed:<a href="#">35538033</a>). Known ligands include EGF, TGFA/TGF- alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF (PubMed:<a href="#">12297049</a>, PubMed:<a href="#">15611079</a>, PubMed:<a href="#">17909029</a>, PubMed:<a href="#">20837704</a>, PubMed:<a href="#">27153536</a>, PubMed:<a href="#">2790960</a>, PubMed:<a href="#">7679104</a>, PubMed:<a href="#">8144591</a>, PubMed:<a href="#">9419975</a>). Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:<a href="#">27153536</a>). May also activate the NF-kappa-B signaling cascade (PubMed:<a href="#">11116146</a>). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:<a href="#">11602604</a>). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:<a href="#">11483589</a>). Positively regulates cell migration via interaction with</p> |

CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:[20462955](#)). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity).

### Cellular Location


Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein Golgi apparatus membrane; Single-pass type I membrane protein. Nucleus membrane; Single-pass type I membrane protein. Endosome. Endosome membrane. Nucleus. Note=In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER (PubMed:17909029, PubMed:20674546). Endocytosed upon activation by ligand (PubMed:17182860, PubMed:17909029, PubMed:27153536, PubMed:2790960). Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-associated fibroblasts (CAF) (PubMed:20551055)

### Tissue Location

Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.