

# eIF1A Rabbit mAb

Catalog # AP75385

## Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	<a href="#">P47813</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	16460

## Additional Information

Gene ID	1964
Other Names	EIF1AX
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.





## Protein Information

Name	EIF1AX
Synonyms	EIF1A, EIF4C
Function	<p>Component of the 43S pre-initiation complex (43S PIC), which binds to the mRNA cap-proximal region, scans mRNA 5'-untranslated region, and locates the initiation codon (PubMed:<a href="#">9732867</a>). This protein enhances formation of the cap-proximal complex (PubMed:<a href="#">9732867</a>). Together with EIF1, facilitates scanning, start codon recognition, promotion of the assembly of 48S complex at the initiation codon (43S PIC becomes 48S PIC after the start codon is reached), and dissociation of aberrant complexes (PubMed:<a href="#">9732867</a>). After start codon location, together with EIF5B orients the initiator methionine-tRNA in a conformation that allows 60S ribosomal subunit joining to form the 80S initiation complex (PubMed:<a href="#">35732735</a>). Is released after 80S initiation complex formation, just after GTP hydrolysis by EIF5B, and before release of EIF5B (PubMed:<a href="#">35732735</a>). Its globular part is located in the A site of the 40S ribosomal subunit (PubMed:<a href="#">35732735</a>). Its interaction with EIF5 during scanning contribute to the maintenance of EIF1 within the open 43S PIC (PubMed:<a href="#">24319994</a>). In contrast to yeast orthologs, does not bind EIF1 (PubMed:<a href="#">24319994</a>).</p>

Cellular Location

Cytoplasm.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.