10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ELK1 Rabbit mAb

Catalog # AP75394

Product Information

Application	WB, IHC-P, FC
Primary Accession	P19419
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	44888

Additional Information

Gene ID	2002
Other Names	ELK1
Dilution	WB~~1:500-1:1000 IHC-P~~N/A FC~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	ELK1 (HGNC:3321)
Function	Transcription factor that binds to purine-rich DNA sequences (PubMed:10799319, PubMed:7889942). Forms a ternary complex with SRF and the ETS and SRF motifs of the serum response element (SRE) on the promoter region of immediate early genes such as FOS and IER2 (PubMed:1630903). Induces target gene transcription upon JNK and MAPK-signaling pathways stimulation (PubMed:7889942).
Cellular Location	Nucleus.
Tissue Location	Lung and testis.

Background


Transcription factor that binds to purine-rich DNA sequences (PubMed:10799319, PubMed:7889942). Forms

a ternary complex with SRF and the ETS and SRF motifs of the serum response element (SRE) on the promoter region of immediate early genes such as FOS and IER2 (PubMed:1630903). Induces target gene transcription upon JNK and MAPK-signaling pathways stimulation (PubMed:7889942).


Images




Western blot analysis of ELK1 in Hela lysates using ELK1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using ELK1 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.