

KDM2A Rabbit mAb

Catalog # AP75429

Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	<u>Q9Y2K7</u>
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	132793

Additional Information

Gene ID	22992
Other Names	KDM2A
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	KDM2A
Function	Histone demethylase that specifically demethylates 'Lys-36' of histone H3, thereby playing a central role in histone code. Preferentially demethylates dimethylated H3 'Lys-36' residue while it has weak or no activity for mono- and tri-methylated H3 'Lys-36'. May also recognize and bind to some phosphorylated proteins and promote their ubiquitination and degradation. Required to maintain the heterochromatic state. Associates with centromeres and represses transcription of small non-coding RNAs that are encoded by the clusters of satellite repeats at the centromere. Required to sustain centromeric integrity and genomic stability, particularly during mitosis. Regulates circadian gene expression by repressing the transcriptional activator activity of CLOCK-BMAL1 heterodimer and RORA in a catalytically-independent manner (PubMed: <u>26037310</u>).
Cellular Location	Nucleus, nucleoplasm. Chromosome Note=Punctate expression throughout the nucleoplasm and enriched in the perinucleolar region (PubMed:19001877, PubMed:20417597). Specifically nucleates at CpG islands where it's presence results in chromatin depleted in H3K36me2 (PubMed:19001877, PubMed:20417597)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.