

FGF1 Rabbit mAb

Catalog # AP75434

Product Information

Application WB Primary Accession P05230

Reactivity Human, Mouse, Rat

Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 17460

Additional Information

Gene ID 2246

Other Names FGF1

Dilution WB~~1/500-1/1000

Format Liquid

Protein Information

Name FGF1

Synonyms FGFA

Function Plays an important role in the regulation of cell survival, cell division,

angiogenesis, cell differentiation and cell migration. Functions as a potent mitogen in vitro. Acts as a ligand for FGFR1 and integrins. Binds to FGFR1 in the presence of heparin leading to FGFR1 dimerization and activation via sequential autophosphorylation on tyrosine residues which act as docking sites for interacting proteins, leading to the activation of several signaling cascades. Binds to integrin ITGAV:ITGB3. Its binding to integrin, subsequent ternary complex formation with integrin and FGFR1, and the recruitment of PTPN11 to the complex are essential for FGF1 signaling. Induces the phosphorylation and activation of FGFR1, FRS2, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:18441324, PubMed:20422052). Can induce angiogenesis

(PubMed: 23469107).


Cellular Location Secreted. Cytoplasm. Cytoplasm, cell cortex. Cytoplasm, cytosol. Nucleus.


Note=Lacks a cleavable signal sequence Within the cytoplasm, it is transported to the cell membrane and then secreted by a non-classical pathway that requires Cu(2+) ions and S100A13. Secreted in a complex with SYT1 (By similarity). Binding of exogenous FGF1 to FGFR facilitates endocytosis followed by translocation of FGF1 across endosomal membrane into the cytosol Nuclear import from the cytosol requires the classical nuclear import

Tissue Location

Predominantly expressed in kidney and brain. Detected at much lower levels in heart and skeletal muscle

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.