10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



FOXC1 Rabbit mAb

Catalog # AP75450

Product Information

ApplicationWB, IPPrimary AccessionQ12948ReactivityMouseHostRabbit

Clonality Monoclonal Antibody

Calculated MW 56789

Additional Information

Gene ID 2296

Other Names FOXC1

Dilution WB~~1/500-1/1000 IP~~N/A

Format Liquid

Protein Information

Name FOXC1

Synonyms FKHL7, FREAC3

Function DNA-binding transcriptional factor that plays a role in a broad range of

cellular and developmental processes such as eye, bones, cardiovascular,

kidney and skin development (PubMed: 11782474, PubMed: 14506133,

PubMed:<u>14578375</u>, PubMed:<u>15277473</u>, PubMed:<u>15299087</u>, PubMed:<u>15684392</u>, PubMed:<u>16449236</u>, PubMed:<u>16492674</u>, PubMed:<u>17210863</u>, PubMed:<u>19279310</u>, PubMed:<u>19793056</u>,

PubMed:<u>25786029</u>, PubMed:<u>27804176</u>, PubMed:<u>27907090</u>). Acts either as a transcriptional activator or repressor (PubMed:<u>11782474</u>). Binds to the consensus binding site 5'- [G/C][A/T]AAA[T/C]AA[A/C]-3' in promoter of target

genes (PubMed:<u>11782474</u>, PubMed:<u>12533514</u>, PubMed:<u>14506133</u>, PubMed:<u>19793056</u>, PubMed:<u>27804176</u>, PubMed:<u>7957066</u>). Upon

DNA-binding, promotes DNA bending (PubMed: 14506133, PubMed: 7957066). Acts as a transcriptional coactivator (PubMed: 26565916). Stimulates Indian

hedgehog (Ihh)-induced target gene expression mediated by the transcription factor GLI2, and hence regulates endochondral ossification (By similarity). Also acts as a transcriptional coregulator by increasing DNA-binding capacity of GLI2 in breast cancer cells (PubMed:26565916). Regulates FOXO1 through binding to a conserved element, 5'-GTAAACAAA-3' in its promoter region, implicating FOXC1 as an important regulator of cell viability and resistance to oxidative stress in the eye (PubMed:17993506). Cooperates with transcription

factor FOXC2 in regulating expression of genes that maintain podocyte integrity (By similarity). Promotes cell growth inhibition by stopping the cell cycle in the G1 phase through TGFB1- mediated signals (PubMed: 12408963). Involved in epithelial-mesenchymal transition (EMT) induction by increasing cell proliferation, migration and invasion (PubMed:20406990, PubMed:22991501). Involved in chemokine CXCL12-induced endothelial cell migration through the control of CXCR4 expression (By similarity). Plays a role in the gene regulatory network essential for epidermal keratinocyte terminal differentiation (PubMed: <u>27907090</u>). Essential developmental transcriptional factor required for mesoderm-derived tissues, such as the somites, skin, bone and cartilage. Positively regulates CXCL12 and stem cell factor expression in bone marrow mesenchymal progenitor cells, and hence plays a role in the development and maintenance of mesenchymal niches for haematopoietic stem and progenitor cells (HSPC). Plays a role in corneal transparency by preventing both blood vessel and lymphatic vessel growth during embryonic development in a VEGF-dependent manner. Involved in chemokine CXCL12-induced endothelial cell migration through the control of CXCR4 expression (By similarity). May function as a tumor suppressor (PubMed:12408963).


Cellular Location

Nucleus Note=Colocalizes with PITX2 isoform 3 in the nucleus at subnuclear chromatine regions (PubMed:16449236). Colocalizes with CBX5 to a heterochromatin-rich region of the nucleus (PubMed:15684392) Colocalizes with GLI2 in the nucleus (By similarity) {ECO:0000250 | UniProtKB:Q61572, ECO:0000269 | PubMed:15684392, ECO:0000269 | PubMed:16449236}

Tissue Location

Expressed in keratinocytes of epidermis and hair follicle (PubMed:27907090). Expressed strongly in microvascular invasion (MVI) formation, basal-like breast cancer (BLBC) and hepatocellular tumors (PubMed:20406990, PubMed:22991501). Expressed in breast cancers (at protein level) (PubMed:26565916). Expressed in hematopoietic cells (PubMed:8499623).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.