

# G3BP Rabbit mAb

Catalog # AP75467

## Product Information

---

|                   |                           |
|-------------------|---------------------------|
| Application       | WB, IHC-P, IHC-F, IP, ICC |
| Primary Accession | <a href="#">Q13283</a>    |
| Reactivity        | Human, Mouse, Rat         |
| Host              | Rabbit                    |
| Clonality         | Monoclonal Antibody       |
| Calculated MW     | 52164                     |

## Additional Information

---

|             |   |
|-------------|---|
| Gene ID     | 10146   |
| Other Names | G3BP1   |
| Dilution    | WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A |
| Format      | Liquid  |

## Protein Information

---

|      |  |
|------|--|
| Name | G3BP1 {ECO:0000303 PubMed:23279204, ECO:0000312 HGNC:HGNC:30292} |
|------|--|

|          |  |
|----------|--|
| Function | <p>Protein involved in various processes, such as stress granule formation and innate immunity (PubMed:<a href="#">12642610</a>, PubMed:<a href="#">20180778</a>, PubMed:<a href="#">23279204</a>, PubMed:<a href="#">30510222</a>, PubMed:<a href="#">30804210</a>). Plays an essential role in stress granule formation (PubMed:<a href="#">12642610</a>, PubMed:<a href="#">20180778</a>, PubMed:<a href="#">23279204</a>, PubMed:<a href="#">32302570</a>, PubMed:<a href="#">32302571</a>, PubMed:<a href="#">32302572</a>, PubMed:<a href="#">34739333</a>, PubMed:<a href="#">35977029</a>, PubMed:<a href="#">36183834</a>, PubMed:<a href="#">36279435</a>, PubMed:<a href="#">36692217</a>, PubMed:<a href="#">37379838</a>). Stress granules are membraneless compartments that store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed:<a href="#">12642610</a>, PubMed:<a href="#">20180778</a>, PubMed:<a href="#">23279204</a>, PubMed:<a href="#">27022092</a>, PubMed:<a href="#">32302570</a>, PubMed:<a href="#">32302571</a>, PubMed:<a href="#">32302572</a>, PubMed:<a href="#">36279435</a>, PubMed:<a href="#">37379838</a>). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA-binding; functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (PubMed:<a href="#">32302570</a>, PubMed:<a href="#">32302571</a>, PubMed:<a href="#">32302572</a>, PubMed:<a href="#">34739333</a>, PubMed:<a href="#">36279435</a>, PubMed:<a href="#">36692217</a>). Also acts as an ATP- and magnesium-dependent helicase: unwinds DNA/DNA, RNA/DNA, and RNA/RNA substrates with comparable efficiency (PubMed:<a href="#">9889278</a>). Acts unidirectionally by moving in the 5' to 3' direction along the bound</p> |
|----------|--|

single-stranded DNA (PubMed:[9889278](#)). Unwinds preferentially partial DNA and RNA duplexes having a 17 bp annealed portion and either a hanging 3' tail or hanging tails at both 5'- and 3'-ends (PubMed:[9889278](#)). Plays an essential role in innate immunity by promoting CGAS and RIGI activity (PubMed:[30510222](#), PubMed:[30804210](#)). Participates in the DNA-triggered cGAS/STING pathway by promoting the DNA binding and activation of CGAS (PubMed:[30510222](#)). Triggers the condensation of cGAS, a process probably linked to the formation of membrane-less organelles (PubMed:[34779554](#)). Also enhances RIGI-induced type I interferon production probably by helping RIGI at sensing pathogenic RNA (PubMed:[30804210](#)). May also act as a phosphorylation- dependent sequence-specific endoribonuclease in vitro: Cleaves exclusively between cytosine and adenine and cleaves MYC mRNA preferentially at the 3'-UTR (PubMed:[11604510](#)).



## Cellular Location


Cytoplasm, cytosol. Perikaryon {ECO:0000250 | UniProtKB:P97855}. Cytoplasm, Stress granule. Nucleus Note=Cytoplasmic in proliferating cells (PubMed:11604510). Cytosolic and partially nuclear in resting cells (PubMed:11604510). Recruited to stress granules in response to arsenite treatment (PubMed:12642610, PubMed:20180778). The unphosphorylated form is recruited to stress granules (PubMed:12642610). HRAS signaling contributes to this process by regulating G3BP dephosphorylation (PubMed:12642610)

## Tissue Location

Ubiquitous..

## Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.