10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Glutathione Peroxidase 4 Rabbit mAb

Catalog # AP75497

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	P36969
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	22175

Additional Information

Gene ID	2879
Other Names	GPX4
Dilution	WB~~1:1000-1:10000 IHC-P~~N/A IHC-F~~N/A
Format	1xPBS(pH 7.4), 150mM NaCl, 50% Glycerol, 0.02% Sodium azide and 0.05% BSA
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	GPX4 {ECO:0000303|PubMed:9705830, ECO:0000312|HGNC:HGNC:4556}
Function	Essential antioxidant peroxidase that directly reduces phospholipid hydroperoxide even if they are incorporated in membranes and lipoproteins (PubMed:40281343). Can also reduce cholesterol hydroperoxide and thymine hydroperoxide (By similarity). Plays a key role in protecting cells from oxidative damage by preventing membrane lipid peroxidation (PubMed:40281343). Required to prevent cells from ferroptosis, a non-apoptotic cell death resulting from an iron- dependent accumulation of lipid reactive oxygen species (PubMed:24439385, PubMed:40281343). The presence of selenocysteine (Sec) versus Cys at the active site is essential for life: it provides resistance to overoxidation and prevents cells against ferroptosis (By similarity). The presence of Sec at the active site is also essential for the survival of a specific type of parvalbumin-positive interneurons, thereby preventing against fatal epileptic seizures (By similarity). May be required to protect cells from the toxicity of ingested lipid hydroperoxides (By similarity). Required for normal sperm development and


	male fertility (By similarity). Essential for maturation and survival of photoreceptor cells (By similarity). Plays a role in a primary T-cell response to viral and parasitic infection by protecting T-cells from ferroptosis and by supporting T-cell expansion (By similarity). Plays a role of glutathione peroxidase in platelets in the arachidonic acid metabolism (PubMed:11115402). Reduces hydroperoxy ester lipids formed by a 15-lipoxygenase that may play a role as down- regulator of the cellular 15-lipoxygenase pathway (By similarity). Can reduce fatty acid-derived hydroperoxides (PubMed:11115402, PubMed:36608588). Can also reduce small soluble hydroperoxides such as H2O2, cumene hydroperoxide and tert-butyl hydroperoxide (PubMed:17630701, PubMed:36608588).
Cellular Location	[Isoform Mitochondrial]: Mitochondrion {ECO:0000250|UniProtKB:O70325}
Tissue Location	Present primarily in testis. Expressed in platelets (at protein level) (PubMed:11115402).

Background


Protects cells against membrane lipid peroxidation and cell death. Required for normal sperm development and male fertility. Could play a major role in protecting mammals from the toxicity of ingested lipid hydroperoxides. Essential for embryonic development. Protects from radiation and oxidative damage.

Images





Western blot analysis of GPX4 in Hela, C6, MCF-7, SKOV-3 lysates using Glutathione Peroxidase 4 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.