

GPR37 Rabbit mAb

Catalog # AP75512

Product Information

Application	WB
Primary Accession	O15354
Reactivity	Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	67114

Additional Information

Gene ID	2861
Other Names	GPR37
Dilution	WB~~1/500-1/1000
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	GPR37
Function	<p>G-protein-coupled receptor that plays a role in several physiological pathways such as resolution of inflammatory pain and oligodendrocyte differentiation (By similarity). Acts as a receptor for several ligands including prosaposin, osteocalcin or neuroprotectin D1. Ligand binding induces endocytosis, followed by an ERK phosphorylation cascade (PubMed:11439185, PubMed:23690594). Acts as a receptor for osteocalcin (OCN) to regulate oligodendrocyte differentiation and central nervous system myelination. Mechanistically, plays a negative role in oligodendrocyte differentiation and myelination during development via activation of the ERK1/2 signaling pathway. Therefore, regulates the stability of myelin or resistance of myelin itself to demyelination. Upon activation by neuroprotectin D1 (NPD1), promotes the activation of phagocytosis in macrophages as well as the shift in cytokine release toward an anti-inflammatory profile, and thus helps to reverse inflammatory pain. In addition, the increased macrophage phagocytosis mediates protection against sepsis upon pathogen infection. Additionally, extracellular vesicles derived from efferocyte express prosaposin, which binds to macrophage GPR37 to increase expression of the efferocytosis receptor TIM4 via an ERK-AP1-dependent signaling axis, leading</p>

to increased macrophage efferocytosis efficiency and accelerated resolution of inflammation (By similarity). May also act as a maturation factor of LRP6, protecting LRP6 from the endoplasmic reticulum (ER)-associated protein degradation (ERAD) and thereby promoting the Wnt/beta-catenin signaling pathway (PubMed:[28341812](#)).


Cellular Location

Cell projection, dendrite. Synapse Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Expressed in brain and spinal cord, and at lower levels in testis, placenta and liver, but no detectable expression observed in any other tissue. When overexpressed in cells, tends to become insoluble and unfolded. Accumulation of the unfolded protein may lead to dopaminergic neuronal death in juvenile Parkinson disease (PDJ).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.