

HADHA Rabbit mAb

Catalog # AP75524

Product Information

Application WB, IHC-P, IHC-F, IP, ICC

Primary Accession P40939

Reactivity Human, Mouse, Rat

Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 83000

Additional Information

Gene ID 3030

Other Names HADHA

Dilution WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A

Format Liquid

Protein Information

Name HADHA

Synonyms HADH

Function Mitochondrial trifunctional enzyme catalyzes the last three of the four

reactions of the mitochondrial beta-oxidation pathway (PubMed: 1550553,

PubMed:<u>29915090</u>, PubMed:<u>30850536</u>, PubMed:<u>8135828</u>,

 $\label{pubMed:31604922} \textbf{PubMed:} \underline{\textbf{31604922}} \textbf{). The mitochondrial beta-oxidation pathway is the major}$

energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA

(PubMed:<u>29915090</u>). Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids

(PubMed: <u>30850536</u>, PubMed: <u>31604922</u>). Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA described here carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities while the trifunctional enzyme subunit beta/HADHB bears the 3-ketoacyl-CoA thiolase

activity (PubMed:<u>29915090</u>, PubMed:<u>30850536</u>, PubMed:<u>8135828</u>). Independently of subunit beta, HADHA also exhibits a cardiolipin

acyltransferase activity that participates in cardiolipin remodeling; cardiolipin is a major mitochondrial membrane phospholipid (PubMed:23152787,

PubMed: <u>31604922</u>). HADHA may act downstream of Tafazzin/TAZ, that remodels monolysocardiolipin (MLCL) to a cardiolipin intermediate, and then


HADHA may continue to remodel this species into mature


tetralinoleoyl-cardiolipin (PubMed:31604922). Has also been proposed to act directly on MLCL; capable of acylating MLCL using different acyl-CoA substrates, with highest activity for oleoyl-CoA (PubMed:23152787).

Cellular Location

Mitochondrion. Mitochondrion inner membrane Note=Protein stability and association with mitochondrion inner membrane do not require HADHB.

Images







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.