

Histone H2A.X Rabbit mAb

Catalog # AP75542

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	P16104
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	15145

Additional Information

Gene ID	3014
Other Names	H2AX
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A FC~~1:20 IP~~1:20-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	H2AX (HGNC:4739)
Function	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.
Cellular Location	Nucleus. Chromosome


Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.


Images




Immunocytochemistry analysis of Histone H2A.X (green) in HL-60 using Histone H2A.X antibody, and DAPI (blue).



Western blot analysis of Histone H2A.X in PC-12 lysates using Histone H2A.X antibody.



Western blot analysis of Histone H2A.X in 3T3, HeLa lysates using Histone H2A.X antibody



Immunohistochemistry analysis of paraffin-embedded Human colon cancer using Histone H2A.X antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.