

# HLA C Rabbit mAb

Catalog # AP75551

## **Product Information**

ApplicationWB, IPPrimary AccessionP10321

Reactivity Human, Mouse

**Host** Rabbit

**Clonality** Monoclonal Antibody

Calculated MW 40649

#### **Additional Information**

**Gene ID** 3107

Other Names HLA-C

**Dilution** WB~~1/500-1/1000 IP~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

## **Protein Information**

Name HLA-C ( HGNC:4933)

Synonyms HLAC

**Function** Antigen-presenting major histocompatibility complex class I (MHCI)

molecule with an important role in reproduction and antiviral immunity

(PubMed:<u>11172028</u>, PubMed:<u>20104487</u>, PubMed:<u>20439706</u>, PubMed:<u>20972337</u>, PubMed:<u>24091323</u>, PubMed:<u>28649982</u>,

PubMed:29312307). In complex with B2M/beta 2 microglobulin displays a restricted repertoire of self and viral peptides and acts as a dominant ligand for inhibitory and activating killer immunoglobulin receptors (KIRs) expressed on NK cells (PubMed:16141329). In an allogeneic setting, such as during pregnancy, mediates interaction of extravillous trophoblasts with KIR on uterine NK cells and regulate trophoblast invasion necessary for placentation and overall fetal growth (PubMed:20972337, PubMed:24091323). During viral infection, may present viral peptides with low affinity for KIRs, impeding KIR-mediated inhibition through peptide antagonism and favoring lysis of infected cells (PubMed:20439706). Presents a restricted repertoire of viral peptides on antigen-presenting cells for recognition by alpha-beta T cell receptor (TCR) on HLA-C-restricted CD8-positive T cells, guiding

antigen-specific T cell immune response to eliminate infected cells, particularly in chronic viral infection settings such as HIV-1 or CMV infection (PubMed:11172028, PubMed:20104487, PubMed:28649982). Both the peptide and the MHC molecule are recognized by TCR, the peptide is responsible for the fine specificity of antigen recognition and MHC residues account for the MHC restriction of T cells (By similarity). Typically presents intracellular peptide antigens of 9 amino acids that arise from cytosolic proteolysis via proteasome. Can bind different peptides containing allele-specific binding motifs, which are mainly defined by anchor residues at position 2 and 9. Preferentially displays peptides having a restricted repertoire of hydrophobic or aromatic amino acids (Phe, Ile, Leu, Met, Val and Tyr) at the C-terminal anchor (PubMed:25311805, PubMed:8265661).


**Cellular Location** 

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

**Tissue Location** 

Ubiquitous. Highly expressed in fetal extravillous trophoblasts in the decidua basalis (at protein level)

# **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.