10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# DYRK1A Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7555c

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Primary Accession** | Q13627 |
| **Reactivity** | Human, Rat, Mouse |
| **Predicted** | Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB12637 |
| **Calculated MW** | 85584 |
| **Antigen Region** | 389-418 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 1859 |
| **Other Names** | Dual specificity tyrosine-phosphorylation-regulated kinase 1A, Dual specificity YAK1-related kinase, HP86, Protein kinase minibrain homolog, MNBH, hMNB, DYRK1A, DYRK, MNB, MNBH |
| **Target/Specificity** | This DYRK1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 389-418 amino acids from the Central region of human DYRK1A. |
| **Dilution** | WB~~1:1000 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | DYRK1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | DYRK1A {ECO:0000303\|PubMed:25620562, ECO:0000312\|HGNC:HGNC:3091} |
| **Function** | Dual-specificity kinase which possesses both serine/threonine and tyrosine |

kinase activities (PubMed:20981014, PubMed:21127067, PubMed:23665168, PubMed:30773093, PubMed:8769099). Exhibits a substrate preference for proline at position P+1 and arginine at position P-3 (PubMed:23665168). Plays an important role in double-strand breaks (DSBs) repair following DNA damage (PubMed:31024071). Mechanistically, phosphorylates RNF169 and increases its ability to block accumulation of TP53BP1 at the DSB sites thereby promoting homologous recombination repair (HRR) (PubMed:30773093). Also acts as a positive regulator of transcription by acting as a CTD kinase that mediates phosphorylation of the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP II) POLR2A (PubMed:25620562, PubMed:29849146). May play a role in a signaling pathway regulating nuclear functions of cell proliferation (PubMed:14500717). Modulates alternative splicing by phosphorylating the splice factor SRSF6 (By similarity). Has pro-survival function and negatively regulates the apoptotic process (By similarity). Promotes cell survival upon genotoxic stress through phosphorylation of SIRT1 (By similarity). This in turn inhibits p53/TP53 activity and apoptosis (By similarity). Phosphorylates SEPTIN4, SEPTIN5 and SF3B1 at 'Thr-434' (By similarity).

**Cellular Location** Nucleus. Nucleus speckle {ECO:0000250|UniProtKB:Q61214}

**Tissue Location** Ubiquitous. Highest levels in skeletal muscle, testis, fetal lung and fetal kidney.


## Background

DYRK1A is a member of the Dual-specificity tyrosine phosphorylation-regulated kinase (DYRK) family. This member contains a nuclear targeting signal sequence, a protein kinase domain, a leucine zipper motif, and a highly conservative 13-consecutive-histidine repeat. It catalyzes its autophosphorylation on serine/threonine and tyrosine residues. It may play a significant role in a signaling pathway regulating cell proliferation and may be involved in brain development. This gene is a homolog of Drosophila mnb (minibrain) gene and rat Dyrk gene. It is localized in the Down syndrome critical region of chromosome 21, and is considered to be a strong candidate gene for learning defects associated with Down syndrome.

## References


Adayev,T., Biochemistry 46 (25), 7614-7624 (2007)
Chang,H.S., Int. J. Cancer 120 (11), 2377-2385 (2007)
Alvarez,M., Mol. Biol. Cell 18 (4), 1167-1178 (2007)
Wissing,J., Mol. Cell Proteomics 6 (3), 537-547 (2007)


## Images




Anti-DYRK1A Antibody (Center) at 1:2000 dilution + NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 86 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.