

IKK gamma Rabbit mAb

Catalog # AP75609

Product Information

Application	WB, IP, ICC
Primary Accession	Q9Y6K9
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	48198

Additional Information




Gene ID	8517
Other Names	IKBKG
Dilution	WB~~1/500-1/1000 IP~~1/20 ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	IKBKG (HGNC:5961)
Synonyms	FIP3, NEMO
Function	Regulatory subunit of the IKK core complex which phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor (PubMed: 14695475 , PubMed: 20724660 , PubMed: 21518757 , PubMed: 9751060). Its binding to scaffolding polyubiquitin plays a key role in IKK activation by multiple signaling receptor pathways (PubMed: 16547522 , PubMed: 18287044 , PubMed: 19033441 , PubMed: 19185524 , PubMed: 21606507 , PubMed: 27777308 , PubMed: 33567255). Can recognize and bind both 'Lys-63'-linked and linear polyubiquitin upon cell stimulation, with a much higher affinity for linear polyubiquitin (PubMed: 16547522 , PubMed: 18287044 , PubMed: 19033441 , PubMed: 19185524 , PubMed: 21606507 , PubMed: 27777308). Could be implicated in NF-kappa-B-mediated protection from cytokine toxicity. Essential for viral activation of IRF3 (PubMed: 19854139). Involved in TLR3- and IFIH1-mediated antiviral innate response; this function requires 'Lys- 27'-linked polyubiquitination (PubMed: 20724660).

Cellular Location	Cytoplasm. Nucleus Note=Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress.
Tissue Location	Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.