

# ITPA Rabbit mAb

Catalog # AP75620

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">Q9BY32</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Calculated MW</b>	21446

## Additional Information

---

<b>Gene ID</b>	3704
<b>Other Names</b>	ITPA
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A
<b>Format</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.



## Protein Information

---

<b>Name</b>	ITPA {ECO:0000255   HAMAP-Rule:MF_03148}
<b>Synonyms</b>	C20orf37
<b>Function</b>	Pyrophosphatase that hydrolyzes the non-canonical purine nucleotides inosine triphosphate (ITP), deoxyinosine triphosphate (dITP) as well as 2'-deoxy-N-6-hydroxylaminopurine triphosphate (dHAPTP) and xanthosine 5'-triphosphate (XTP) to their respective monophosphate derivatives. The enzyme does not distinguish between the deoxy- and ribose forms. Probably excludes non-canonical purines from RNA and DNA precursor pools, thus preventing their incorporation into RNA and DNA and avoiding chromosomal lesions.
<b>Cellular Location</b>	Cytoplasm {ECO:0000255   HAMAP-Rule:MF_03148, ECO:0000269   PubMed:11278832}
<b>Tissue Location</b>	Ubiquitous. Highly expressed in heart, liver, sex glands, thyroid and adrenal gland

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.