10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Protein Jumonji Rabbit mAb

Catalog # AP75637

Product Information

Application	WB
Primary Accession	Q92833
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	138734

Additional Information

Gene ID	3720
Other Names	JARID2
Dilution	WB~~1:1000-1:5000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	JARID2
Synonyms	JMJ
Function	Regulator of histone methyltransferase complexes that plays an essential role in embryonic development, including heart and liver development, neural tube fusion process and hematopoiesis (PubMed:20075857). Acts as an accessory subunit for the core PRC2 (Polycomb repressive complex 2) complex, which mediates histone H3K27 (H3K27me3) trimethylation on chromatin (PubMed:20075857, PubMed:29499137, PubMed:31959557). Binds DNA and mediates the recruitment of the PRC2 complex to target genes in embryonic stem cells, thereby playing a key role in stem cell differentiation and normal embryonic development (PubMed:20075857). In cardiac cells, it is required to repress expression of cyclin-D1 (CCND1) by activating methylation of 'Lys-9' of histone H3 (H3K9me) by the GLP1/EHMT1 and G9a/EHMT2 histone methyltransferases (By similarity). Also acts as a transcriptional repressor of ANF via its interaction with GATA4 and NKX2-5 (By similarity).

	Participates in the negative regulation of cell proliferation signaling (By similarity). Does not have histone demethylase activity (By similarity).
Cellular Location	Nucleus {ECO:0000255|PROSITE-ProRule:PRU00355, ECO:0000255|PROSITE-ProRule:PRU00537, ECO:0000269|PubMed:20075857, ECO:0000269|PubMed:29499137}. Note=Colocalizes with the PRC2 complex on chromatin.
Tissue Location	During embryogenesis, predominantly expressed in neurons and particularly in dorsal root ganglion cells

Background

Regulator of histone methyltransferase complexes that plays an essential role in embryonic development, including heart and liver development, neural tube fusion process and hematopoiesis.

Images



Western blot analysis of Jarid2 in K562, C6 lysates using Protein Jumonji antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.