

Junctional Adhesion Molecule 1/JAM-A Rabbit mAb

Catalog # AP75641

Product Information

Application	WB, IHC-P
Primary Accession	Q9Y624
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	32583

Additional Information

Gene ID	50848
Other Names	F11R
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	F11R
Synonyms	JAM1, JCAM
Function	Seems to play a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3 (PubMed: 11489913). The association of the PARD6-PARD3 complex may prevent the interaction of PARD3 with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier (By similarity). Ligand for integrin alpha-L/beta-2 involved in memory T- cell and neutrophil transmigration (PubMed: 11812992). Involved in platelet activation (PubMed: 10753840).
Cellular Location	Cell junction, tight junction. Cell membrane; Single-pass type I membrane protein. Note=Localized at tight junctions of both epithelial and endothelial cells.


Tissue Location

Expressed in endothelium, epithelium and leukocytes (at protein level).


Background

Seems to play a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3. The association of the PARD6-PARD3 complex may prevent the interaction of PARD3 with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier. Involved in platelet activation. In case of orthoreovirus infection, serves as receptor for the virus.

Images



Western blot analysis of Junctional Adhesion Molecule 1/JAMA in K562, PC-3 lysates using Junctional Adhesion Molecule 1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Junctional Adhesion Molecule 1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.