

# LXR alpha Rabbit mAb

Catalog # AP75684

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P              |
| Primary Accession | <a href="#">Q13133</a> |
| Reactivity        | Human, Rat             |
| Host              | Rabbit                 |
| Clonality         | Monoclonal Antibody    |
| Calculated MW     | 50396                  |

## Additional Information

---

|             |  |
|-------------|--|
| Gene ID     | 10062  |
| Other Names | NR1H3  |
| Dilution    | WB~~1/500-1/1000 IHC-P~~N/A  |
| Format      | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.    |
| Storage     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

## Protein Information

---

|          |  |
|----------|--|
| Name     | NR1H3  |
| Synonyms | LXRA   |
| Function | <p>Nuclear receptor that exhibits a ligand-dependent transcriptional activation activity (PubMed:<a href="#">19481530</a>, PubMed:<a href="#">25661920</a>, PubMed:<a href="#">37478846</a>).</p> <p>Interaction with retinoic acid receptor (RXR) shifts RXR from its role as a silent DNA-binding partner to an active ligand- binding subunit in mediating retinoid responses through target genes defined by LXRES (PubMed:<a href="#">37478846</a>). LXRES are DR4-type response elements characterized by direct repeats of two similar hexanuclotide half-sites spaced by four nucleotides (By similarity).</p> <p>Plays an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8 (PubMed:<a href="#">19481530</a>). Interplays functionally with RORA for the regulation of genes involved in liver metabolism (By similarity).</p> <p>Induces LPCAT3-dependent phospholipid remodeling in endoplasmic reticulum (ER) membranes of hepatocytes, driving SREBF1 processing and lipogenesis (By similarity). Via LPCAT3, triggers the incorporation of arachidonate into phosphatidylcholines of ER membranes, increasing membrane dynamics and enabling triacylglycerols transfer to nascent very</p> |

low-density lipoprotein (VLDL) particles. Via LPCAT3 also counteracts lipid-induced ER stress response and inflammation, likely by modulating SRC kinase membrane compartmentalization and limiting the synthesis of lipid inflammatory mediators (By similarity).

## Cellular Location




Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00407, ECO:0000269 | PubMed:25661920}. Cytoplasm {ECO:0000250 | UniProtKB:Q9Z0Y9}

## Tissue Location

Visceral organs specific expression. Strong expression was found in liver, kidney and intestine followed by spleen and to a lesser extent the adrenals

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.