

mH2A1 Rabbit mAb

Catalog # AP75686

Product Information

Application	WB, IHC-P
Primary Accession	O75367
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	39184

Additional Information

Gene ID	9555
Other Names	MACROH2A1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	MACROH2A1 (HGNC:4740)
Function	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes where it represses transcription (PubMed: 12718888 , PubMed: 15621527 , PubMed: 16428466). Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template (PubMed: 15897469). Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability (PubMed: 15897469). DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Involved in stable X chromosome inactivation (PubMed: 15897469). Inhibits the binding of transcription factors, including NF-kappa-B, and interferes with the activity of remodeling SWI/SNF complexes (PubMed: 12718888 , PubMed: 16428466). Inhibits histone acetylation by EP300 and recruits class I HDACs, which induces a hypoacetylated state of chromatin (PubMed: 16107708 , PubMed: 16428466).

Cellular Location


Nucleus. Chromosome. Note=Enriched in inactive X chromosome chromatin and in senescence-associated heterochromatin (PubMed:15621527, PubMed:15897469, PubMed:9634239). Recruited to DNA damage sites in an APLF-dependent manner (PubMed:21211722, PubMed:29905837).

Tissue Location


Widely expressed..

Background

Involved in stable X chromosome inactivation. Inhibits the binding of transcription factors and interferes with the activity of remodeling SWI/SNF complexes. Inhibits histone acetylation by EP300 and recruits class I HDACs, which induces an hypoacetylated state of chromatin.

Images

Western blot analysis of macroH2A.1 in rat Brain, C6, 3T3 lysates using Macro H2A.1 antibody.



Immunocytochemistry analysis of macroH2A.1(green) in Hela using macroH2A.1 antibody,and DAPI(blue)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.