

MAGE3 Rabbit mAb

Catalog # AP75688

Product Information

Application	WB, IP
Primary Accession	P43357
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	34747

Additional Information

Gene ID	4102
Other Names	MAGEA3
Dilution	WB~~1:500-1:1000 IP~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	MAGEA3 {ECO:0000303 PubMed:29779948, ECO:0000312 HGNC:HGNC:6801}
Function	Activator of ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases that acts as a repressor of autophagy (PubMed: 20864041 , PubMed: 31267705). May enhance ubiquitin ligase activity of TRIM28 and stimulate p53/TP53 ubiquitination by TRIM28. Proposed to act through recruitment and/or stabilization of the Ubl-conjugating enzyme (E2) at the E3:substrate complex (PubMed: 17942928 , PubMed: 20864041). May play a role in embryonal development and tumor transformation or aspects of tumor progression (PubMed: 17942928 , PubMed: 20864041). In vitro promotes cell viability in melanoma cell lines (PubMed: 17942928). Antigen recognized on a melanoma by autologous cytolytic T-lymphocytes (PubMed: 8113684).
Tissue Location	Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not

in normal tissues except for testes and placenta. Never expressed in kidney tumors, Leukemias and lymphomas

Background
