

PUM2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7569b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q8TB72
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10534
Calculated MW	114216
Antigen Region	133-162

Additional Information

Gene ID	23369
Other Names	Pumilio homolog 2, Pumilio-2, PUM2, KIAA0235, PUMH2
Target/Specificity	This PUM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 133-162 amino acids from the N-terminal region of human PUM2.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PUM2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PUM2
Synonyms	KIAA0235, PUMH2
Function	Sequence-specific RNA-binding protein that acts as a post- transcriptional repressor by binding the 3'-UTR of mRNA targets. Binds to an RNA consensus sequence, the Pumilio Response Element (PRE), 5'- UGUANAUA-3', that is

related to the Nanos Response Element (NRE) ([PubMed:21397187](#)). Mediates post-transcriptional repression of transcripts via different mechanisms: acts via direct recruitment of the CCR4-POP2-NOT deadenylase leading to translational inhibition and mRNA degradation ([PubMed:22955276](#)). Also mediates deadenylation-independent repression by promoting accessibility of miRNAs ([PubMed:18776931](#), [PubMed:22345517](#)). Acts as a post-transcriptional repressor of E2F3 mRNAs by binding to its 3'-UTR and facilitating miRNA regulation ([PubMed:22345517](#)). Plays a role in cytoplasmic sensing of viral infection ([PubMed:25340845](#)). Represses a program of genes necessary to maintain genomic stability such as key mitotic, DNA repair and DNA replication factors. Its ability to repress those target mRNAs is regulated by the lncRNA NORAD (non-coding RNA activated by DNA damage) which, due to its high abundance and multitude of PUMILIO binding sites, is able to sequester a significant fraction of PUM1 and PUM2 in the cytoplasm ([PubMed:26724866](#)). May regulate DCUN1D3 mRNA levels ([PubMed:25349211](#)). May support proliferation and self-renewal of stem cells. Binds specifically to miRNA MIR199A precursor, with PUM1, regulates miRNA MIR199A expression at a postranscriptional level ([PubMed:28431233](#)).

Cellular Location

Cytoplasm. Cytoplasmic granule. Cytoplasm, perinuclear region. Note=The cytoplasmic granules are stress granules which are a dense aggregation in the cytosol composed of proteins and RNAs that appear when the cell is under stress. Colocalizes with NANOS3 in the stress granules Colocalizes with NANOS1 and SNAPIN in the perinuclear region of germ cells.

Tissue Location

Expressed in male germ cells of adult testis (at protein level). Highly expressed in testis and ovary. Predominantly expressed in stem cells and germ cells. Expressed at lower level in brain, heart, kidney, liver, muscle, placenta, intestine and stomach Expressed in cerebellum, corpus callosum, caudate nucleus, hippocampus, medulla oblongata and putamen. Expressed in all fetal tissues tested

Background



PUM2 is a member of the PUF family, evolutionarily conserved RNA-binding proteins related to the Pumilio proteins of *Drosophila* and the fem-3 mRNA binding factor proteins of *C. elegans*. This protein contains a sequence-specific RNA binding domain comprised of eight repeats and N- and C-terminal flanking regions, and serves as a translational regulator of specific mRNAs by binding to their 3' untranslated regions. The evolutionarily conserved function of this protein in invertebrates and lower vertebrates suggests that the human protein may be involved in translational regulation of embryogenesis, and cell development and differentiation.

References

Gupta,Y.K., Structure 16 (4), 549-557 (2008)
 Spassov,D.S.,IUBMB Life 55 (7), 359-366 (2003)
 Spassov,D.S.,Gene 299 (1-2), 195-204 (2002)

Images

Western blot analysis of PUM2 (arrow) using rabbit polyclonal PUM2 Antibody (N-term) (Cat.#AP7569b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PUM2 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human testis tissue reacted with PUM2 Antibody (C-term) (Cat.#AP7569b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.