

Nogo B Receptor Rabbit mAb

Catalog # AP75815

Product Information

Application	WB
Primary Accession	Q96E22
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	33224

Additional Information

Gene ID	116150
Other Names	NUS1
Dilution	WB~~1:1000-1:5000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	NUS1 {ECO:0000303 PubMed:28842490, ECO:0000312 HGNC:HGNC:21042}
Function	With DHDDS, forms the dehydrodolichyl diphosphate synthase (DDS) complex, an essential component of the dolichol monophosphate (Dol-P) biosynthetic machinery (PubMed: 21572394 , PubMed: 25066056 , PubMed: 28842490 , PubMed: 32817466 , PubMed: 33077723). Both subunits contribute to enzymatic activity, i.e. condensation of multiple copies of isopentenyl pyrophosphate (IPP) to farnesyl pyrophosphate (FPP) to produce dehydrodolichyl diphosphate (Dedol-PP), a precursor of dolichol phosphate which is utilized as a sugar carrier in protein glycosylation in the endoplasmic reticulum (ER) (PubMed: 21572394 , PubMed: 25066056 , PubMed: 28842490 , PubMed: 32817466 , PubMed: 33077723). Synthesizes long- chain polyprenols, mostly of C95 and C100 chain length (PubMed: 32817466). Regulates the glycosylation and stability of nascent NPC2, thereby promoting trafficking of LDL-derived cholesterol (PubMed: 21572394). Acts as a specific receptor for the N-terminus of Nogo-B, a neural and cardiovascular regulator (PubMed: 16835300).


Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein.
Note=Colocalizes with Nogo-B during VEGF and wound healing angiogenesis

Background

With DHDDS, forms the dehydrodolichyl diphosphate synthase (DDS) complex, an essential component of the dolichol monophosphate (Dol-P) biosynthetic machinery. Both subunits contribute to enzymatic activity, i.e. condensation of multiple copies of isopentenyl pyrophosphate (IPP) to farnesyl pyrophosphate (FPP) to produce dehydrodolichyl diphosphate (Dedol-PP), a precursor of dolichol phosphate which is utilized as a sugar carrier in protein glycosylation in the endoplasmic reticulum (ER) (PubMed:21572394, PubMed:25066056, PubMed:28842490). Regulates the glycosylation and stability of nascent NPC2, thereby promoting trafficking of LDL-derived cholesterol. Acts as a specific receptor for the N-terminus of Nogo-B, a neural and cardiovascular regulator (PubMed:16835300).

Images



Western blot analysis of Nogo B receptor in K562, 3T3, HeLa lysates using Nogo B Receptor antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.