

NUR77 Rabbit mAb

Catalog # AP75835

Product Information

Application	WB
Primary Accession	P22736
Reactivity	Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	64463


Additional Information

Gene ID	3164
Other Names	NR4A1
Dilution	WB~~1/500-1/1000
Format	Liquid

Protein Information

Name	NR4A1
Synonyms	GFRP1, HMR, NAK1
Function	<p>Orphan nuclear receptor. Binds the NGFI-B response element (NBRE) 5'-AAAGGTCA-3' (PubMed:18690216, PubMed:8121493, PubMed:9315652). Binds 9-cis-retinoic acid outside of its ligand- binding (NR LBD) domain (PubMed:18690216). Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation (PubMed:22983157). Regulates the inflammatory response in macrophages by regulating metabolic adaptations during inflammation, including repressing the transcription of genes involved in the citric acid cycle (TCA) (By similarity). Inhibits NF-kappa-B signaling by binding to low-affinity NF-kappa-B binding sites, such as at the IL2 promoter (PubMed:15466594). May act concomitantly with NR4A2 in regulating the expression of delayed-early genes during liver regeneration (By similarity). Plays a role in the vascular response to injury (By similarity).</p>
Cellular Location	<p>Nucleus. Cytoplasm, cytosol. Mitochondrion Note=Nuclear export to the cytosol is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340). Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950).</p>
Tissue Location	Fetal muscle and adult liver, brain and thyroid.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.