

PABPN1 Rabbit mAb

Catalog # AP75861

Product Information

Application	WB, IHC-P, IHC-F, FC, IP
Primary Accession	Q86U42
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	32749

Additional Information

Gene ID	8106
Other Names	PABPN1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IHC-F~~N/A FC~~1:20-1:50 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	PABPN1 (HGNC:8565)
Synonyms	PAB2, PABP2
Function	Involved in the 3'-end formation of mRNA precursors (pre- mRNA) by the addition of a poly(A) tail of 200-250 nt to the upstream cleavage product (By similarity). Stimulates poly(A) polymerase (PAPOLA) conferring processivity on the poly(A) tail elongation reaction and also controls the poly(A) tail length (By similarity). Increases the affinity of poly(A) polymerase for RNA (By similarity). Is also present at various stages of mRNA metabolism including nucleocytoplasmic trafficking and nonsense-mediated decay (NMD) of mRNA. Cooperates with SKIP to synergistically activate E-box-mediated transcription through MYOD1 and may regulate the expression of muscle- specific genes (PubMed: 11371506). Binds to poly(A) and to poly(G) with high affinity (By similarity). May protect the poly(A) tail from degradation (By similarity). Subunit of the trimeric poly(A) tail exosome targeting (PAXT) complex, a complex that directs a subset of long and polyadenylated poly(A) RNAs for

exosomal degradation. The RNA exosome is fundamental for the degradation of RNA in eukaryotic nuclei. Substrate targeting is facilitated by its cofactor MTREX, which links to RNA-binding protein adapters (PubMed:[27871484](#)).

Cellular Location

Nucleus. Cytoplasm. Nucleus speckle Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Shuttles between the nucleus and the cytoplasm but predominantly found in the nucleus (PubMed:10688363). Its nuclear import may involve the nucleocytoplasmic transport receptor transportin and a RAN-GTP- sensitive import mechanism (By similarity). Is exported to the cytoplasm by a carrier-mediated pathway that is independent of mRNA traffic. Colocalizes with SKIP and poly(A) RNA in nuclear speckles (By similarity). Intranuclear filamentous inclusions or 'aggregates' are detected in the myocytes of patients; these inclusions contain PABPN1, ubiquitin, subunits of the proteasome and poly(A) RNA {ECO:0000250|UniProtKB:Q28165, ECO:0000269|PubMed:10688363, ECO:0000269|PubMed:11001936, ECO:0000269|PubMed:11371506, ECO:0000269|PubMed:14663186, ECO:0000269|PubMed:17289661, ECO:0000269|PubMed:27209344}


Tissue Location

Ubiquitous.

Background





This gene encodes an abundant nuclear protein that binds with high affinity to nascent poly(A) tails. The protein is required for progressive and efficient polymerization of poly(A) tails at the 3' ends of eukaryotic transcripts and controls the size of the poly(A) tail to about 250 nt. At steady-state, this protein is localized in the nucleus whereas a different poly(A) binding protein is localized in the cytoplasm. This gene contains a GCG trinucleotide repeat at the 5' end of the coding region, and expansion of this repeat from the normal 6 copies to 8-13 copies leads to autosomal dominant oculopharyngeal muscular dystrophy (OPMD) disease. Related pseudogenes have been identified on chromosomes 19 and X. Read-through transcription also exists between this gene and the neighboring upstream BCL2-like 2 (BCL2L2) gene.

Images



Western blot analysis of PABPN1 in Jurkat, HeLa, COS7, Raji lysates using PABPN1 antibody.

Immunohistochemistry analysis of paraffin-embedded Human breast cancer using PABPN1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.