

PBK Rabbit mAb

Catalog # AP75873

Product Information

Application	WB, IHC-P, IP
Primary Accession	Q96KB5
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	36085

Additional Information

Gene ID	55872
Other Names	PBK
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	PBK
Synonyms	TOPK
Function	Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage.
Tissue Location	Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.

Background


This genes encodes a serine/threonine kinase related to the dual specific mitogen-activated protein kinase

kinase (MAPKK) family. Evidence suggests that mitotic phosphorylation is required for its catalytic activity. This mitotic kinase may be involved in the activation of lymphoid cells and support Ticular functions, with a suggested role in the process of spermatogenesis.


Images



Immunohistochemistry analysis of paraffin-embedded mouse colon using PBK antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of PBK in K562, HeLa lysates using PBK antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.