

PDK2 Rabbit mAb

Catalog # AP75884

Product Information

Application WB, IHC-P, IP **Primary Accession** 015119

Reactivity Human, Mouse, Rat

Host Rabbi

Clonality Monoclonal Antibody

Calculated MW 46154

Additional Information

Gene ID 5164

Other Names PDK2

Dilution WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name PDK2

Synonyms PDHK2

Function Kinase that plays a key role in the regulation of glucose and fatty acid

dehydrogenase subunits PDHA1 and PDHA2. This inhibits pyruvate dehydrogenase activity, and thereby regulates metabolite flux through the tricarboxylic acid cycle, down-regulates aerobic respiration and inhibits the formation of acetyl-coenzyme A from pyruvate. Inhibition of pyruvate dehydrogenase decreases glucose utilization and increases fat metabolism. Mediates cellular responses to insulin. Plays an important role in maintaining normal blood glucose levels and in metabolic adaptation to nutrient

metabolism and homeostasis via phosphorylation of the pyruvate

availability. Via its regulation of pyruvate dehydrogenase activity, plays an important role in maintaining normal blood pH and in preventing the accumulation of ketone bodies under starvation. Plays a role in the regulation of cell proliferation and in resistance to apoptosis under oxidative stress.


Plays a role in p53/TP53-mediated apoptosis.

Cellular Location Mitochondrion matrix.

Tissue Location

Expressed in many tissues, with the highest level in heart and skeletal muscle, intermediate levels in brain, kidney, pancreas and liver, and low levels in placenta and lung

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.