

[KD]PDLIM7 Rabbit mAb

Catalog # AP75886

Product Information

Application	WB, IHC-P, FC
Primary Accession	Q9NR12
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	49845

Additional Information

Gene ID	9260
Other Names	PDLIM7
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A FC~~1:50-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	PDLIM7
Synonyms	ENIGMA
Function	May function as a scaffold on which the coordinated assembly of proteins can occur. May play a role as an adapter that, via its PDZ domain, localizes LIM-binding proteins to actin filaments of both skeletal muscle and nonmuscle tissues. Involved in both of the two fundamental mechanisms of bone formation, direct bone formation (e.g. embryonic flat bones mandible and cranium), and endochondral bone formation (e.g. embryonic long bone development). Plays a role during fracture repair. Involved in BMP6 signaling pathway (By similarity).
Cellular Location	Cytoplasm. Cytoplasm, cytoskeleton. Note=Colocalizes with RET to the cell periphery and in some cytoskeletal components. Colocalizes with TPM2 near the Z line in muscle. Colocalizes with TBX4 and TBX5 to actin filaments (By similarity).


Tissue Location

Isoform 1 and isoform 2 are expressed ubiquitously, however, isoform 2 predominates in skeletal muscle, isoform 1 is more abundant in lung, spleen, leukocytes and fetal liver

Background

The protein encoded by this gene is representative of a family of proteins composed of conserved PDZ and LIM domains. LIM domains are proposed to function in protein-protein recognition in a variety of contexts including gene transcription and development and in cytoskeletal interaction. The LIM domains of this protein bind to protein kinases, whereas the PDZ domain binds to actin filaments. The gene product is involved in the assembly of an actin filament-associated complex essential for transmission of ret/ptc2 mitogenic signaling. The biological function is likely to be that of an adapter, with the PDZ domain localizing the LIM-binding proteins to actin filaments of both skeletal muscle and nonmuscle tissues. Alternative splicing of this gene results in multiple transcript variants.

Images



Western blot analysis of PDLIM7 in HeLa lysates using PDLIM7 antibody.

