

PEPD Rabbit mAb

Catalog # AP75891

Product Information

Application	WB
Primary Accession	P12955
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	54548

Additional Information

Gene ID	5184
Other Names	PEPD
Dilution	WB~~1:500-1:1000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	PEPD {ECO:0000303 PubMed:8198124, ECO:0000312 HGNC:HGNC:8840}
Function	Dipeptidase that catalyzes the hydrolysis of dipeptides with a prolyl (Xaa-Pro) or hydroxyprolyl residue in the C-terminal position (PubMed: 17081196 , PubMed: 35165443). The preferred dipeptide substrate is Gly-Pro, but other Xaa-Pro dipeptides, such as Ala-Pro, Met-Pro, Phe-Pro, Val-Pro and Leu-Pro, can be cleaved (PubMed: 17081196). Plays an important role in collagen metabolism because the high level of iminoacids in collagen (PubMed: 2925654).

Background

Splits dipeptides with a prolyl or hydroxyprolyl residue in the C-terminal position. Plays an important role in collagen metabolism because the high level of iminoacids in collagen.

Images



Western blot analysis of PEPD in K562, HeLa lysates using PEPD antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.