

# Cullin 4A (CUL4A) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7590a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q13619</a>
Other Accession	<a href="#">Q3TCH7</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB10557
Calculated MW	87680
Antigen Region	102-131

## Additional Information

---

Gene ID	8451
Other Names	Cullin-4A, CUL-4A, CUL4A
Target/Specificity	This Cullin 4A (CUL4A) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 102-131 amino acids from the N-terminal region of human Cullin 4A (CUL4A).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Cullin 4A (CUL4A) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	CUL4A {ECO:0000303   PubMed:9721878, ECO:0000312   HGNC:HGNC:2554}
Function	Core component of multiple cullin-RING-based E3 ubiquitin- protein ligase complexes which mediate the ubiquitination of target proteins (PubMed: <a href="#">14578910</a> , PubMed: <a href="#">14739464</a> , PubMed: <a href="#">15448697</a> ,

PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#), PubMed:[30166453](#), PubMed:[33854232](#), PubMed:[33854239](#)). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DET1-COP1) directs ubiquitination of JUN (PubMed:[14739464](#)). DCX(DDB2) directs ubiquitination of XPC (PubMed:[15811626](#)). DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition (PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of p53/TP53 in response to radiation-induced DNA damage and during DNA replication (PubMed:[14578910](#), PubMed:[15448697](#), PubMed:[15548678](#)). DCX(DTL) directs autoubiquitination of DTL (PubMed:[23478445](#)). In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip (PubMed:[16537899](#)). Is involved in ubiquitination of HOXA9 (PubMed:[14609952](#)). The DDB1-CUL4A- DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed:[26431207](#)). The DCX(ERCC8) complex (also named CSA complex) plays a role in transcription-coupled repair (TCR) (PubMed:[12732143](#), PubMed:[32355176](#), PubMed:[38316879](#)). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:[29779948](#)). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiquitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:[33854232](#), PubMed:[33854239](#)). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed:[30166453](#)). With CUL4B, contributes to ribosome biogenesis (PubMed:[26711351](#)).

## Background

---

CUL4a is a core component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1. The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component. DCX(DET1-COP1) directs ubiquitination of JUN. DCX(DDB2) directs ubiquitination of XPC. In association with RBX1, DDB1 and DDB2 is required for histone H3 and histone H4 ubiquitination in response to ultraviolet and may be important for subsequent DNA repair. DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of TP53 in response to

radiation-induced DNA damage and during DNA replication. In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip. Is involved in ubiquitination of HOXA9.


## References

---

Huang,J.,Oncogene 27 (29), 4056-4064 (2008)  
Hu,J.,Genes Dev. 22 (7), 866-871 (2008)  
Wen,X., J. Biol. Chem. 282 (37), 27046-27057 (2007)

## Images

---



Western blot analysis of CUL4a (arrow) using rabbit polyclonal CUL4a Antibody (Human N-term) (Cat.#AP7590a).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CUL4a gene (Lane 2) (Origene Technologies).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.