

PKN1 Rabbit mAb

Catalog # AP75925

Product Information

Application	WB
Primary Accession	Q16512
Reactivity	Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	103932

Additional Information

Gene ID	5585
Other Names	PKN1
Dilution	WB~~1/500-1/1000
Format	Liquid

Protein Information


Name	PKN1
Synonyms	PAK1, PKN, PRK1, PRKCL1
Function	<p>PKC-related serine/threonine-protein kinase involved in various processes such as regulation of the intermediate filaments of the actin cytoskeleton, cell migration, tumor cell invasion and transcription regulation. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14. Regulates the cytoskeletal network by phosphorylating proteins such as VIM and neurofilament proteins NEFH, NEFL and NEFM, leading to inhibit their polymerization. Phosphorylates 'Ser-575', 'Ser-637' and 'Ser-669' of MAPT/Tau, lowering its ability to bind to microtubules, resulting in disruption of tubulin assembly. Acts as a key coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and specifically mediating phosphorylation of 'Thr-11' of histone H3 (H3T11ph), a specific tag for epigenetic transcriptional activation that promotes demethylation of histone H3 'Lys-9' (H3K9me) by KDM4C/JMJD2C. Phosphorylates HDAC5, HDAC7 and HDAC9, leading to impair their import in the nucleus. Phosphorylates 'Thr-38' of PPP1R14A, 'Ser-159', 'Ser-163' and 'Ser-170' of MARCKS, and GFAP. Able to phosphorylate RPS6 in vitro.</p>
Cellular Location	Cytoplasm. Nucleus Endosome. Cell membrane {ECO:0000250 UniProtKB:Q63433}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q63433}. Cleavage furrow. Midbody
Note=Associates with chromatin in a ligand-dependent manner Localization
to endosomes is mediated via its interaction with RHOB Association to the cell
membrane is dependent on Ser-377 phosphorylation. Accumulates during
telophase at the cleavage furrow and finally concentrates around the
midbody in cytokinesis {ECO:0000250|UniProtKB:Q63433,
ECO:0000269|PubMed:17332740}

Tissue Location

Found ubiquitously. Expressed in heart, brain, placenta, lung, skeletal muscle,
kidney and pancreas. Expressed in numerous tumor cell lines, especially in
breast tumor cells

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.