

PMCA1 Rabbit mAb

Catalog # AP75929

Product Information

Application	WB, IHC-P, IP
Primary Accession	P20020
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	134685

Additional Information

Gene ID	490
Other Names	ATP2B1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ATP2B1 (HGNC:814)
Function	Catalyzes the hydrolysis of ATP coupled with the transport of calcium from the cytoplasm to the extracellular space thereby maintaining intracellular calcium homeostasis (PubMed: 35358416). Plays a role in blood pressure regulation through regulation of intracellular calcium concentration and nitric oxide production leading to regulation of vascular smooth muscle cells vasoconstriction. Positively regulates bone mineralization through absorption of calcium from the intestine. Plays dual roles in osteoclast differentiation and survival by regulating RANKL-induced calcium oscillations in preosteoclasts and mediating calcium extrusion in mature osteoclasts (By similarity). Regulates insulin sensitivity through calcium/calmodulin signaling pathway by regulating AKT1 activation and NOS3 activation in endothelial cells (PubMed: 29104511). May play a role in synaptic transmission by modulating calcium and proton dynamics at the synaptic vesicles.
Cellular Location	Cell membrane; Multi-pass membrane protein. Basolateral cell membrane

{ECO:0000250|UniProtKB:G5E829}. Synapse
{ECO:0000250|UniProtKB:G5E829} Presynaptic cell membrane
{ECO:0000250|UniProtKB:G5E829}; Multi-pass membrane protein.
Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane
{ECO:0000250|UniProtKB:G5E829}; Multi-pass membrane protein.
Note=Colocalizes with SV2A in photoreceptor synaptic terminals. Colocalizes with NPTN to the immunological synapse. Colocalizes with EPB41 to the basolateral membrane in enterocyte. Preferentially sorted to recycling synaptic vesicles. {ECO:0000250|UniProtKB:G5E829}


Tissue Location

Isoform B: Ubiquitously expressed. Isoform C: Found in brain cortex, skeletal muscle and heart muscle. Isoform D: Has only been found in fetal skeletal muscle. Isoform K: Found in small intestine and liver. Abundantly expressed in the endometrial epithelial cells and glandular epithelial cells in early-proliferative phase and early-secretory phases (PubMed:21400627)


Background

The protein encoded by this gene belongs to the family of P-type primary ion transport ATPases characterized by the formation of an aspartyl phosphate intermediate during the reaction cycle. These enzymes remove bivalent calcium ions from eukaryotic cells against very large concentration gradients and play a critical role in intracellular calcium homeostasis. The mammalian plasma membrane calcium ATPase isoforms are encoded by at least four separate genes and the diversity of these enzymes is further increased by alternative splicing of transcripts. The expression of different isoforms and splice variants is regulated in a developmental, tissue- and cell type-specific manner, suggesting that these pumps are functionally adapted to the physiological needs of particular cells and tissues. This gene encodes the plasma membrane calcium ATPase isoform 1. Alternatively spliced transcript variants encoding different isoforms have been identified.


Images



Western blot analysis of PMCA1 in mouse brain, rat lung lysates using PMCA1 antibody.



Western blot analysis of PMCA1 in U87-MG, Human placenta lysates using PMCA1 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.