

[KO]PRAME Rabbit mAb

Catalog # AP75943

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	P78395
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	57890

Additional Information

Gene ID	23532
Other Names	PRAME
Dilution	WB~~1:1000-1:10000 IHC-P~~N/A FC~~1:200-1:500 IP~~1:20-1:50
Format	1xPBS(pH 7.4), 150mM NaCl, 50% Glycerol, 0.02% Sodium azide and 0.05% BSA
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	PRAME {ECO:0000303 PubMed:9047241, ECO:0000312 HGNC:HGNC:9336}
Function	Substrate-recognition component of a Cul2-RING (CRL2) E3 ubiquitin-protein ligase complex, which mediates ubiquitination of target proteins, leading to their degradation (PubMed: 21822215 , PubMed: 26138980). The CRL2(PRAME) complex mediates ubiquitination and degradation of truncated MSRB1/SEPX1 selenoproteins produced by failed UGA/Sec decoding (PubMed: 26138980). In the nucleus, the CRL2(PRAME) complex is recruited to epigenetically and transcriptionally active promoter regions bound by nuclear transcription factor Y (NFY) and probably plays a role in chromatin regulation (PubMed: 21822215). Functions as a transcriptional repressor, inhibiting the signaling of retinoic acid through the retinoic acid receptors RARA, RARB and RARG: prevents retinoic acid-induced cell proliferation arrest, differentiation and apoptosis (PubMed: 16179254).
Cellular Location	Nucleus. Chromosome. Cytoplasm Golgi apparatus. Cell membrane. Note=Associates with chromatin; specifically enriched at transcriptionally

active promoters that are also bound by nuclear transcription factor Y (composed of NFYA, NFYB and NFYC) and at enhancers (PubMed:21822215). Recruited to the Golgi apparatus in response to interferon gamma (IFNG) treatment (PubMed:23460923).


Tissue Location

Expressed in testis. Detected in samples of kidney, brain and skin.


Background

Substrate-recognition component of a Cul2-RING (CRL2) E3 ubiquitin-protein ligase complex, which mediates ubiquitination of target proteins, leading to their degradation (PubMed:21822215, PubMed:26138980).


Images





Western blot analysis of PRAME in HCT116(WT) and PRAME-Knockout HCT116(KO) lysates using PRAME antibody.



Immunohistochemistry analysis of paraffin-embedded mouse testis using PRAME antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of PRAME in K562 lysates using PRAME antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.