10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

[KD]Rab5 Rabbit mAb

Catalog # AP75986

Product Information

Application	WB, IHC-P, IHC-F, FC, IP
Primary Accession	P20339
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	23659

Additional Information

Gene ID	5868
Other Names	RAB5A
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IHC-F~~N/A FC~~1:20-1:50 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	RAB5A (HGNC:9783)
Synonyms	RAB5
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes and involved in early endocytic trafficking (PubMed:10818110, PubMed:14617813, PubMed:15378032, PubMed:16086013, PubMed:16410077, PubMed:17562788). Required for EEA1 recruitment to early endosomes (PubMed:16086013, PubMed:17562788). Recruits FERRY complex to early endosomes, where FERRY links early endosomes with a subgroup of mRNAs to enable mRNA intracellular distribution (PubMed:37267906). Recruits RABEP1/Rabaptin- 5

	effector to early endosomes, thereby promoting endocytic membrane fusion (By similarity). Required for EGF and transferrin endocytosis and trafficking through early endosomes (PubMed:16086013, PubMed:17562788). Contributes to the regulation of filopodia extension (PubMed:14978216). Required for the exosomal release of SDCBP, CD63, PDCD6IP and syndecan (PubMed:22660413). Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 (By similarity).
Cellular Location	Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane; Lipid-anchor. Melanosome Cytoplasmic vesicle. Cell projection, ruffle {ECO:0000250|UniProtKB:P18066}. Membrane. Cytoplasm, cytosol. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9CQD1}. Endosome membrane Note=Enriched in stage I melanosomes (PubMed:17081065). Alternates between membrane-bound and cytosolic forms (Probable) {ECO:0000269|PubMed:17081065, ECO:0000305}


Background

Rab5 is a member of the Ras superfamily of small Rab GTPases. Rab5 is localized at the plasma membrane and early endosomes and functions as a key regulator of vesicular trafficking during early endocytosis (1). The conformational change between Rab5 GTP/GDP states is essential for its biological function as a rate limiting regulator at multiple steps during endocytosis.

Images





Western blot analysis of Rab5 in HeLa(WT) and Rab5-Knockdown HeLa(KD) using Rab5 antibody.




Immunocytochemistry analysis of Rab5 (green) in HT-1080 using Rab5 antibody,and DAPI(blue).

Western blot analysis of Rab5 in K562, C6, 3T3, Hela lysates using Rab5 antibody.







Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Rab5 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.