

# Rab9 Rabbit mAb

Catalog # AP75991

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">P51151</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	22838

## Additional Information

---

<b>Gene ID</b>	9367
<b>Other Names</b>	RAB9A
<b>Dilution</b>	WB~~1:1000-1:5000 IHC-P~~N/A
<b>Format</b>	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	RAB9A ( <a href="#">HGNC:9792</a> )
<b>Synonyms</b>	RAB9
<b>Function</b>	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). RAB9A is involved in the transport of proteins between the endosomes and the trans-Golgi network (TGN) (PubMed: <a href="#">34793709</a> ). Specifically uses NDE1/NDEL1 as an effector to interact with the dynein motor complex in order to control retrograde trafficking of RAB9-associated late endosomes to the TGN (PubMed: <a href="#">34793709</a> ). Involved in the recruitment of SGSM2 to melanosomes and is required for the proper trafficking of melanogenic enzymes TYR, TYRP1 and DCT/TYRP2 to melanosomes in melanocytes (By similarity).


## Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Endoplasmic reticulum membrane. Golgi apparatus membrane. Late endosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle membrane. Melanosome {ECO:0000250|UniProtKB:Q9R0M6}. Note=Colocalizes with OSBPL1A at the late endosome (PubMed:16176980). Recruited to phagosomes containing *S.aureus* or *M.tuberculosis* (PubMed:21255211). Mainly localizes to late endosomes and partially localizes to Golgi (PubMed:34793709) Colocalizes with NDE1 to membrane vesicles (PubMed:34793709)


## Background

Involved in the transport of proteins between the endosomes and the trans Golgi network. Interacts (preferentially in its GTP-bound form) with GCC2 (via its GRIP domain).

## Images



Western blot analysis of Rab9 in 3T3, HeLa lysates using Rab9 antibody.



Immunohistochemistry analysis of paraffin-embedded Human colon cancer using Rab9 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

