

RBPJK Rabbit mAb

Catalog # AP76014

Product Information

ApplicationWBPrimary AccessionQ06330ReactivityRatHostRabbit

Clonality Monoclonal Antibody

Calculated MW 55637

Additional Information

Gene ID 3516

Other Names RBPJ

Dilution WB~~1/500-1/1000

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name RBPJ (HGNC:5724)

Function Transcriptional regulator that plays a central role in Notch signaling, a

signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD

product of Notch proteins (Notch intracellular domain), it acts as a

transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA (PubMed:21991380). Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen)


subunits (By similarity).

Cellular Location Nucleus. Cytoplasm. Note=Mainly nuclear, upon interaction with

RITA/C12orf52, translocates to the cytoplasm, down-regulating the Notch

(PubMed:<u>23303788</u>). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.