

# AXL Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7602a

#### **Product Information**

**Application** WB, IHC-P, FC, E

Primary Accession
Reactivity
Human
Host
Clonality
Polyclonal
Isotype
Rabbit IgG
Calculated MW
98337
Antigen Region
P30530
Human
Rabbit
Rabbit
98337

#### **Additional Information**

Gene ID 558

Other Names Tyrosine-protein kinase receptor UFO, AXL oncogene, AXL, UFO

**Target/Specificity** This AXL antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 13-44 amino acids from the N-terminal

region of human AXL.

**Dilution** WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent

concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** AXL Antibody (N-term) is for research use only and not for use in diagnostic or

therapeutic procedures.

### **Protein Information**

Name AXL

Synonyms UFO

**Function** Receptor tyrosine kinase that transduces signals from the extracellular

matrix into the cytoplasm by binding growth factor GAS6 and which is thus

regulating many physiological processes including cell survival, cell

proliferation, migration and differentiation. Ligand binding at the cell surface induces dimerization and autophosphorylation of AXL. Following activation by ligand, AXL binds and induces tyrosine phosphorylation of PI3-kinase subunits PIK3R1, PIK3R2 and PIK3R3; but also GRB2, PLCG1, LCK and PTPN11. Other downstream substrate candidates for AXL are CBL, NCK2, SOCS1 and TNS2. Recruitment of GRB2 and phosphatidylinositol 3 kinase regulatory subunits by AXL leads to the downstream activation of the AKT kinase. GAS6/AXL signaling plays a role in various processes such as endothelial cell survival during acidification by preventing apoptosis, optimal cytokine signaling during human natural killer cell development, hepatic regeneration, gonadotropin-releasing hormone neuron survival and migration, platelet activation, or regulation of thrombotic responses. Also plays an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response.

**Cellular Location** 

Cell membrane; Single-pass type I membrane protein

**Tissue Location** 

Highly expressed in metastatic colon tumors. Expressed in primary colon tumors. Weakly expressed in normal colon tissue.


## **Background**

AXL, a member of the AXL/UFO subfamily of Tyr protein kinases, may function as a signal transducer between specific cell types of mesodermal origin. This Type I membrane protein has transforming potential in patients with chronic myeloproliferative disorder or chronic myelocytic leukemia. The protein contains 2 putative fibronectin type III domains and 2 putative immunoglobulin-like C2-type domains.

## References


Lee, S.T., et al., Oncogene 8(12):3403-3410 (1993). Janssen, J.W., et al., Oncogene 6(11):2113-2120 (1991). O'Bryan, J.P., et al., Mol. Cell. Biol. 11(10):5016-5031 (1991). Partanen, J., et al., Proc. Natl. Acad. Sci. U.S.A. 87(22):8913-8917 (1990).


# **Images**



Western blot analysis of anti-AXL N-term Pab (Cat. #AP7602a) in SKBR3 cell lysate. AXL (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.





Flow cytometric analysis of WiDr cells using AXL Antibody (N-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.