10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SEC6 Rabbit mAb

Catalog # AP76040

Product Information

Application	WB, IHC-P, IP
Primary Accession	O60645
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	85567

Additional Information

Gene ID	11336
Other Names	EXOC3
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IP~~1:20
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	EXOC3
Synonyms	SEC6, SEC6L1
Function	Component of the exocyst complex involved in the docking of exocytic vesicles with fusion sites on the plasma membrane.
Cellular Location	Cytoplasm {ECO:0000250|UniProtKB:O54921}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:O54921}. Cell projection, growth cone {ECO:0000250|UniProtKB:O54921}. Midbody. Golgi apparatus. Cell projection, neuron projection {ECO:0000250|UniProtKB:Q62825}. Note=Perinuclear in undifferentiated cells. Redistributes to growing neurites and growth cones during neuronal differentiation (By similarity). During mitosis, early recruitment to the midbody requires RALA, but not RALB, and EXOC2. In late stages of cytokinesis, localization to the midbody is RALB- dependent (PubMed:18756269). {ECO:0000250|UniProtKB:O54921, ECO:0000269|PubMed:18756269}

Tissue Location	Expressed in epididymis (at protein level).


Background

Exoc3 (Sec6) is a component of the exocyst (Sec6/8) complex involved in vesicle trafficking at neural synapses and vesicle targeting in polarized epithelial cells. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. This complex is composed of eight proteins (Sec3, Sec5, Sec6, Sec8, Sec10, and Sec15, Exo70 and Exo84), with molecular weights ranging from 70 to 144 kDa.


Images




Western blot analysis of rSec6 in rat Brain, 3T3, Hela lysates using Exocyst Complex Component 3 antibody.



Immunocytochemistry analysis of rSec6(green) in Hela using rSec6 antibody,and DAPI(blue)





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.