10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

LUM Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7605c

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P51884
Other Accession	P51885, Q05443
Reactivity	Human
Predicted	Bovine, Mouse, Rat, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18423
Calculated MW	38429
Antigen Region	198-227

Additional Information

Gene ID	4060
Other Names	Lumican, Keratan sulfate proteoglycan lumican, KSPG lumican, LUM, LDC, SLRR2D
Target/Specificity	This LUM antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 198-227 amino acids from the Central region of human LUM.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LUM Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LUM
Synonyms	LDC, SLRR2D

Cellular Location Secreted, extracellular space, extracellular matrix

Tissue Location Cornea and other tissues.


Background

LUM is a member of the small leucine-rich proteoglycan (SLRP) family that includes decorin, biglycan, fibromodulin, keratocan, epiphycan, and osteoglycin. In these bifunctional molecules, this protein moiety binds collagen fibrils and the highly charged hydrophilic glycosaminoglycans regulate interfibrillar spacings. The protein is the major keratan sulfate proteoglycan of the cornea but is also distributed in interstitial collagenous matrices throughout the body. Lumican may regulate collagen fibril organization and circumferential growth, corneal transparency, and epithelial cell migration and tissue repair.


References

Wang,P., Li,S. Invest. Ophthalmol. Vis. Sci. 50 (4), 1546-1551 (2009)
Matsuda,Y., Yamamoto,T. Int. J. Oncol. 33 (6), 1177-1185 (2008)
Kelemen,L.E., Couch,F.J. Breast Cancer Res. 10 (6), R98 (2008)

Images




Western blot analysis of LUM antibody (Center) (Cat.#AP7605c) in A375 cell line lysates (35ug/lane). LUM (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human prostate carcinoma reacted with LUM Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

LUM Antibody (Center) (Cat. #AP7605c) flow cytometric analysis of A375 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.