10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SCIN Rabbit mAb

Catalog # AP76060

Product Information

Application	WB, IHC-P
Primary Accession	Q9Y6U3
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	80489

Additional Information

Gene ID	85477
Other Names	SCIN
Dilution	WB~~1:500-1:1000 IHC-P~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	SCIN {ECO:0000303|PubMed:8547642, ECO:0000312|HGNC:HGNC:21695}
Function	Ca(2+)-dependent actin filament-severing protein that has a regulatory function in exocytosis by affecting the organization of the microfilament network underneath the plasma membrane (PubMed:26365202, PubMed:8547642). Severing activity is inhibited by phosphatidylinositol 4,5-bis-phosphate (PIP2) (By similarity). In vitro, also has barbed end capping and nucleating activities in the presence of Ca(2+). Required for megakaryocyte differentiation, maturation, polyploidization and apoptosis with the release of platelet-like particles (PubMed:11568009). Plays a role in osteoclastogenesis (OCG) and actin cytoskeletal organization in osteoclasts (By similarity). Regulates chondrocyte proliferation and differentiation (By similarity). Inhibits cell proliferation and tumorigenesis. Signaling is mediated by MAPK, p38 and JNK pathways (PubMed:11568009).
Cellular Location	Cytoplasm, cytoskeleton. Cell projection, podosome {ECO:0000250|UniProtKB:Q60604}

Tissue Location	Expressed in megakaryocytes.


Background

Ca2+-dependent actin filament-severing protein that has a regulatory function in exocytosis by affecting the organization of the microfilament network underneath the plasma membrane (PubMed:8547642, PubMed:26365202). Severing activity is inhibited by phosphatidylinositol 4,5-bis-phosphate (PIP2) . In vitro, also has barbed end capping and nucleating activities in the presence of Ca2+. Required for megakaryocyte differentiation, maturation, polyploidization and apoptosis with the release of platelet-like particles (PubMed:11568009). Plays a role in osteoclastogenesis (OCG) and actin cytoskeletal organization in osteoclasts . Regulates chondrocyte proliferation and differentiation . Inhibits cell proliferation and tumorigenesis. Signaling is mediated by MAPK, p38 and JNK pathways (PubMed:11568009).

Images



Western blot analysis of SCIN in Hela lysates using SCIN antibody.



Immunohistochemistry analysis of paraffin-embedded Human colon cancer using SCIN antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.