

# PLA2G2A Rabbit mAb

Catalog # AP76064

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P14555</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	16083

## Additional Information

---

<b>Gene ID</b>	5320
<b>Other Names</b>	PLA2G2A
<b>Dilution</b>	WB~~1:1000-1:5000
<b>Format</b>	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	PLA2G2A
<b>Synonyms</b>	PLA2B, PLA2L, RASF-A
<b>Function</b>	Secretory calcium-dependent phospholipase A2 that primarily targets extracellular phospholipids with implications in host antimicrobial defense, inflammatory response and tissue regeneration (PubMed: <a href="#">10455175</a> , PubMed: <a href="#">10681567</a> , PubMed: <a href="#">2925633</a> ). Hydrolyzes the ester bond of the fatty acyl group attached at sn-2 position of phospholipids (phospholipase A2 activity) with preference for phosphatidylethanolamines and phosphatidylglycerols over phosphatidylcholines (PubMed: <a href="#">10455175</a> , PubMed: <a href="#">10681567</a> ). Contributes to lipid remodeling of cellular membranes and generation of lipid mediators involved in pathogen clearance. Displays bactericidal activity against Gram-positive bacteria by directly hydrolyzing phospholipids of the bacterial membrane (PubMed: <a href="#">10358193</a> , PubMed: <a href="#">11694541</a> ). Upon sterile inflammation, targets membrane phospholipids of extracellular mitochondria released from activated platelets,

generating free unsaturated fatty acids such as arachidonate that is used by neighboring leukocytes to synthesize inflammatory eicosanoids such as leukotrienes. Simultaneously, by compromising mitochondrial membrane integrity, promotes the release in circulation of potent damage-associated molecular pattern molecules that activate the innate immune response (PubMed:[25082876](#)). Plays a stem cell regulator role in the intestinal crypt. Within intracellular compartment mediates Paneth cell differentiation and its stem cell supporting functions by inhibiting Wnt signaling pathway in intestinal stem cell (ICS). Secreted in the intestinal lumen upon inflammation, acts in an autocrine way and promotes prostaglandin E2 synthesis that stimulates Wnt signaling pathway in ICS cells and tissue regeneration (By similarity). May play a role in the biosynthesis of N-acyl ethanolamines that regulate energy metabolism and inflammation. Hydrolyzes N-acyl phosphatidylethanolamines to N-acyl lysophosphatidylethanolamines, which are further cleaved by a lysophospholipase D to release N-acyl ethanolamines (PubMed:[14998370](#)). Independent of its catalytic activity, acts as a ligand for integrins (PubMed:[18635536](#), PubMed:[25398877](#)). Binds to and activates integrins ITGAV:ITGB3, ITGA4:ITGB1 and ITGA5:ITGB1 (PubMed:[18635536](#), PubMed:[25398877](#)). Binds to a site (site 2) which is distinct from the classical ligand-binding site (site 1) and induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:[25398877](#)). Induces cell proliferation in an integrin-dependent manner (PubMed:[18635536](#)).

#### Cellular Location

Secreted. Cell membrane; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein


#### Tissue Location

Expressed in various tissues including heart, kidney, liver, lung, pancreas, placenta, skeletal muscle, prostate, ovary, colon and small intestine. Not detected in lymphoid organs and brain (PubMed:10455175, PubMed:10681567). Expressed in platelets (at protein level) (PubMed:25082876).

## Background

Thought to participate in the regulation of the phospholipid metabolism in biomembranes including eicosanoid biosynthesis. Catalyzes the calcium-dependent hydrolysis of the 2-acyl groups in 3-sn-phosphoglycerides.

## Images



Western blot analysis of Secretory Phospholipase A2 in K562, C6, 3T3, HeLa lysates using Secretory Phospholipase A2 antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.