10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



SF3B3 Rabbit mAb

Catalog # AP76074

Product Information

Application WB. ICC 015393 **Primary Accession**

Human, Mouse, Rat Reactivity

Host

Clonality Monoclonal Antibody

Calculated MW 135577

Additional Information

Gene ID 23450

Other Names SF3B3

Dilution WB~~1/500-1/1000 ICC~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name SF3B3

Synonyms KIAA0017, SAP130 {ECO:0000303 | PubMed:104

Component of the 17S U2 SnRNP complex of the spliceosome, a large **Function**

ribonucleoprotein complex that removes introns from transcribed pre-mRNAs

(PubMed: 10490618, PubMed: 10882114, PubMed: 12234937, PubMed: <u>27720643</u>, PubMed: <u>28781166</u>, PubMed: <u>32494006</u>,

PubMed:34822310). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA


branch- site adenosine, the nucleophile for the first step of splicing


(PubMed: 12234937, PubMed: 32494006, PubMed: 34822310). Within the 17S U2 SnRNP complex, SF3B3 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed:12234937, PubMed:27720643). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:12234937). May also be involved in the assembly of the 'E' complex (PubMed: 10882114). Also acts as a component of the minor spliceosome,


Cellular Location

Nucleus

Images







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.