

SMAC Rabbit mAb

Catalog # AP76094

Product Information

Application	WB, IHC-P, IHC-F, IP
Primary Accession	Q9NR28
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	27131

Additional Information

Gene ID	56616
Other Names	DIABLO
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IHC-F~~N/A IP~~1:10-1:100
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	DIABLO (HGNC:21528)
Function	Promotes apoptosis by activating caspases in the cytochrome c/Apaf-1/caspase-9 pathway. Acts by opposing the inhibitory activity of inhibitor of apoptosis proteins (IAP). Inhibits the activity of BIRC6/BRUCE by inhibiting its binding to caspases (PubMed: 15200957 , PubMed: 36758104 , PubMed: 36758105 , PubMed: 36758106).
Cellular Location	Mitochondrion. Cytoplasm, cytosol Note=Released into the cytosol in a PARL-dependent manner when cells undergo apoptosis.
Tissue Location	Ubiquitously expressed with highest expression in testis. Expression is also high in heart, liver, kidney, spleen, prostate and ovary. Low in brain, lung, thymus and peripheral blood leukocytes. Isoform 3 is ubiquitously expressed


Background

Smac/Diablo is a 21 kDa mammalian mitochondrial protein that functions as a regulatory component during apoptosis. Upon mitochondrial stress, Smac/Diablo is released from mitochondria and competes with caspases for binding of IAPs (inhibitor of apoptosis proteins). The interaction of Smac/Diablo with IAPs relieves the inhibitory effect of the IAPs on caspases.


Images



Western blot analysis of Smac/Diablo in mouse heart, rat skin lysates using SMAC antibody.




Western blot analysis of Smac/Diablo in K562, HeLa lysates using Smac/Diablo antibody.



Immunocytochemistry analysis of Smac/Diablo(green) in HeLa using Smac/Diablo antibody, and DAPI(blue)

Immunohistochemistry analysis of paraffin-embedded Human tonsil using Smac/Diablo antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.