

STK33 Rabbit mAb

Catalog # AP76128

Product Information

Application	WB, IHC-P, IHC-F
Primary Accession	Q9BYT3
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	57831

Additional Information

Gene ID	65975
Other Names	STK33
Dilution	WB~~1:500-1:1000 IHC-P~~N/A IHC-F~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	STK33 {ECO:0000303 PubMed:34155512}
Function	Serine/threonine protein kinase required for spermatid differentiation and male fertility (PubMed: 37146716 , PubMed: 38781365). Promotes sperm flagella assembly during spermatogenesis by mediating phosphorylation of fibrous sheath proteins AKAP3 and AKAP4 (By similarity). Also phosphorylates vimentin/VIM, thereby regulating the dynamic behavior of the intermediate filament cytoskeleton (By similarity).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q924X7}. Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:Q924X7}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:Q924X7}. Note=Colocalizes with the caudal end of the manchette, a transient structure that guides tail elongation in elongating spermatids {ECO:0000250 UniProtKB:Q924X7}
Tissue Location	Highly expressed in testis, fetal lung and heart, followed by pituitary gland, kidney, interventricular septum, pancreas, heart, trachea, thyroid gland and

uterus. Weak hybridization signals were observed in the following tissues: amygdala, aorta, esophagus, colon ascending, colon transverse, skeletal muscle, spleen, peripheral blood leukocyte, lymph node, bone marrow, placenta, prostate, liver, salivary gland, mammary gland, some tumor cell lines, fetal brain, fetal liver, fetal spleen and fetal thymus. No signal at all was detectable in RNA from tissues of the nervous system



Background

Serine/threonine protein kinase which phosphorylates VIME. May play a specific role in the dynamic behavior of the intermediate filament cytoskeleton by phosphorylation of VIME (By similarity). Not essential for the survival of KRAS-dependent AML cell lines.

Images



Western blot analysis of STK33 in Jurkat, 293 lysates using STK33 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.