

Stromal Interaction Molecule 1 Rabbit mAb

Catalog # AP76130

Product Information

Application	WB, IHC-P
Primary Accession	Q13586
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	77423

Additional Information

Gene ID	6786
Other Names	STIM1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	STIM1
Synonyms	GOK {ECO:0000303 PubMed:9377559}
Function	Acts as a Ca(2+) sensor that gates two major inward rectifying Ca(2+) channels at the plasma membrane: Ca(2+) release- activated Ca(2+) (CRAC) channels and arachidonate-regulated Ca(2+)- selective (ARC) channels (PubMed: 15866891 , PubMed: 16005298 , PubMed: 16208375 , PubMed: 16537481 , PubMed: 16733527 , PubMed: 16766533 , PubMed: 16807233 , PubMed: 18854159 , PubMed: 19182790 , PubMed: 19249086 , PubMed: 19622606 , PubMed: 19706554 , PubMed: 22464749 , PubMed: 24069340 , PubMed: 24351972 , PubMed: 24591628 , PubMed: 25326555 , PubMed: 26322679 , PubMed: 28219928 , PubMed: 32415068). Plays a role in mediating store-operated Ca(2+) entry (SOCE), a Ca(2+) influx following depletion of intracellular Ca(2+) stores. Upon Ca(2+) depletion, translocates from the endoplasmic reticulum to the plasma membrane where it activates CRAC

channel pore-forming subunits ORA1, ORA2 and ORA3 to generate sustained and oscillatory Ca²⁺ entry (PubMed:[16208375](#), PubMed:[16537481](#), PubMed:[32415068](#)). Involved in enamel formation (PubMed:[24621671](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endoplasmic reticulum membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton. Sarcoplasmic reticulum. Note=Translocates from the endoplasmic reticulum to the cell membrane in response to a depletion of intracellular calcium and is detected at punctae corresponding to junctions between the endoplasmic reticulum and the cell membrane (PubMed:[16005298](#), PubMed:[16208375](#), PubMed:[18854159](#), PubMed:[19182790](#), PubMed:[19249086](#)). Associated with the microtubule network at the growing distal tip of microtubules (PubMed:[19632184](#)). Colocalizes with ORAI1 at the cell membrane (PubMed:[27185316](#)). Colocalizes preferentially with CASQ1 at endoplasmic reticulum in response to a depletion of intracellular calcium (PubMed:[27185316](#))


Tissue Location

Ubiquitously expressed in various human primary cells and tumor cell lines.


Background

STIM1 is a potential tumor suppressor; defects in STIM1 may cause rhabdomyosarcoma and rhabdoid tumors. STIM1 can either homodimerize or form heterodimers with STIM2. STIM2 possesses a high sequence identity to STIM1 and can function as an inhibitor of STIM1-mediated plasma membrane store-operated Ca²⁺ entry. However, further investigation is required to elucidate the true physiological function of STIM2.

Images



Western blot analysis of Stromal interaction molecule 1 in MCF-7, C6, 3T3, HeLa lysates using Stromal Interaction Molecule 1 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.