

# TAK1 Rabbit mAb

Catalog # AP76149

## Product Information

---

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	<a href="#">O43318</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	67196

## Additional Information

---

Gene ID	6885
Other Names	MAP3K7
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

Name	MAP3K7 {ECO:0000303   PubMed:28397838, ECO:0000312   HGNC:HGNC:6859}
Function	Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway (PubMed: <a href="#">10094049</a> , PubMed: <a href="#">11460167</a> , PubMed: <a href="#">12589052</a> , PubMed: <a href="#">16845370</a> , PubMed: <a href="#">16893890</a> , PubMed: <a href="#">21512573</a> , PubMed: <a href="#">8663074</a> , PubMed: <a href="#">9079627</a> ). Plays an important role in the cascades of cellular responses evoked by changes in the environment (PubMed: <a href="#">10094049</a> , PubMed: <a href="#">11460167</a> , PubMed: <a href="#">12589052</a> , PubMed: <a href="#">16845370</a> , PubMed: <a href="#">16893890</a> , PubMed: <a href="#">21512573</a> , PubMed: <a href="#">8663074</a> , PubMed: <a href="#">9079627</a> ). Mediates signal transduction of TRAF6, various cytokines including interleukin-1 (IL-1), transforming growth factor- beta (TGFB), TGFB-related factors like BMP2 and BMP4, toll-like receptors (TLR), tumor necrosis factor receptor CD40 and B-cell receptor (BCR) (PubMed: <a href="#">16893890</a> , PubMed: <a href="#">9079627</a> ). Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal transduction cascade through the phosphorylation and activation of several MAP kinase kinases like MAP2K1/MEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7 (PubMed: <a href="#">11460167</a> , PubMed: <a href="#">8663074</a> ). These MAP2Ks in turn activate p38 MAPKs and c-jun N- terminal kinases (JNKs); both p38 MAPK and

JNK pathways control the transcription factors activator protein-1 (AP-1) (PubMed:[11460167](#), PubMed:[12589052](#), PubMed:[8663074](#)). Independently of MAP2Ks and p38 MAPKs, acts as a key activator of NF-kappa-B by promoting activation of the I-kappa-B-kinase (IKK) core complex (PubMed:[12589052](#), PubMed:[8663074](#)). Mechanistically, recruited to polyubiquitin chains of RIPK2 and IKBKG/NEMO via TAB2/MAP3K7IP2 and TAB3/MAP3K7IP3, and catalyzes phosphorylation and activation of IKBKB/IKKB component of the IKK complex, leading to NF-kappa-B activation (PubMed:[10094049](#), PubMed:[11460167](#)). In osmotic stress signaling, plays a major role in the activation of MAPK8/JNK1, but not that of NF-kappa-B (PubMed:[16893890](#)). Promotes TRIM5 capsid-specific restriction activity (PubMed:[21512573](#)). Phosphorylates RIPK1 at 'Ser-321' which positively regulates RIPK1 interaction with RIPK3 to promote necroptosis but negatively regulates RIPK1 kinase activity and its interaction with FADD to mediate apoptosis (By similarity). Phosphorylates STING1 in response to cGAMP-activation, promoting association between STEEP1 and STING1 and STING1 translocation to COPII vesicles (PubMed:[37832545](#)).


### Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Although the majority of MAP3K7/TAK1 is found in the cytosol, when complexed with TAB1/MAP3K7IP1 and TAB2/MAP3K7IP2, it is also localized at the cell membrane

### Tissue Location

Isoform 1A is the most abundant in ovary, skeletal muscle, spleen and blood mononuclear cells. Isoform 1B is highly expressed in brain, kidney and small intestine. Isoform 1C is the major form in prostate. Isoform 1D is the less abundant form

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.